



THE UNIVERSITY *of* EDINBURGH

This thesis has been submitted in fulfilment of the requirements for a postgraduate degree (e.g. PhD, MPhil, DClinPsychol) at the University of Edinburgh. Please note the following terms and conditions of use:

This work is protected by copyright and other intellectual property rights, which are retained by the thesis author, unless otherwise stated.

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge.

This thesis cannot be reproduced or quoted extensively from without first obtaining permission in writing from the author.

The content must not be changed in any way or sold commercially in any format or medium without the formal permission of the author.

When referring to this work, full bibliographic details including the author, title, awarding institution and date of the thesis must be given.

Self-Warmth, Self-Coldness and Self-Esteem as Predictors of Quality of Life and Disability in Chronic Pain

A systematic review of the impact of fibromyalgia on quality of life using the Short Form-36 and an empirical study of the relationship between self-warmth, self-coldness and self-esteem on quality of life and disability in a chronic pain population

Caitríona Doughty



Doctorate in Clinical Psychology

The University of Edinburgh

2016

Table of Contents

Declaration of Own Work	4
Acknowledgements	6
List of Tables	7
List of Figures	7
List of Appendices	8
List of Abbreviations	9
Word Counts	10
Full Thesis Abstract	11
Lay Thesis Summary	13
Chapter 1: A systematic review and meta-analysis of the impact of fibromyalgia on quality of life using the Short Form-36	14
<i>Abstract</i>	<i>15</i>
<i>1.1 Introduction</i>	<i>16</i>
1.1.1. Aims of the Research	18
<i>1.2 Materials and Method</i>	<i>18</i>
1.2.1. Search Strategy	18
1.2.2. Selection Criteria	19
1.2.3. Data Extraction and Quality Assessment	20
1.2.4. Statistical Analysis	21
<i>1.3 Results</i>	<i>21</i>
1.3.1. Search Results	21
1.3.2. Characteristics of Included Studies	22
1.3.3. Methodological Quality Assessment	22
1.3.4. Sample Characteristics	22
1.3.5. Settings	23
1.3.6 Statistical Analysis	23
1.3.7 Meta-analysis of the eight subdomains of the Sf-36	28
1.3.8 Meta-analysis of the two subcomponents of the SF-36 (PCS and MCS)	29
1.3.9 Sensitivity and subgroup analysis	31
1.3.10 Narrative of subdomain findings	31
<i>1.4 Discussion</i>	
1.4.1 Strengths, limitations and implications for future research	37
1.5. References	39

Chapter 2: Self-Warmth, Self-Coldness, and Self-Esteem as Predictors of Quality of Life and Disability in a Chronic Pain Population	47
<i>Abstract</i>	48
<i>2.1 Introduction</i>	49
2.1.1 Evidence for Self-Compassion as a predictor	49
2.1.2 How Self-Compassion is conceptualised and measured	50
2.1.3 Overlap with other concepts	52
2.1.4 Summary	54
2.1.5 Aims	54
<i>2.2 Method</i>	55
2.2.1 Design	55
2.2.2 Participants	56
2.2.3 Measures	56
2.2.4 Procedure	59
2.2.5 Statistical Analysis	59
2.2.6 Power and Sample Size	60
2.2.7 Missing Data	61
2.2.8 Tests of Reliability	61
2.2.9 Data Analysis	62
<i>2.3 Results</i>	62
2.3.1. Descriptive Data	62
2.3.2 Control Variables	64
2.3.3. Correlational Analysis	64
2.3.4. Regression Analysis	64
<i>2.4 Discussion</i>	71
2.4.1 Clinical Implications	76
2.4.2 Strengths and Limitations	76
2.4.3 Conclusions	78
<i>2.5 References</i>	79

DClinPsychol Declaration of Own Work

Name: Caitríona Doughty

Title of Work: Self-Warmth, Self-Coldness and Self-Esteem as Predictors of Quality of Life and Disability in Chronic Pain

I confirm that this work is my own except where indicated, and that I have:

- Read and understood the Plagiarism Rules and Regulations
- Composed and undertaken the work myself
- Clearly referenced/listed all sources as appropriate
- Referenced and put in inverted commas any quoted text of more than three words (from books, web, etc.)
- Given the sources of all pictures, data etc. that are not my own
- Not made undue use of essay(s) of any other student(s), either past or present (or where used, this has been referenced appropriately)
- Not sought or used the help of any external professional agencies for the work (or where used, this has been referenced appropriately)
- Not submitted the work for any other degree or professional qualification except as specified
- Acknowledged in appropriate places any help that I have received from others (e.g. fellow students, technicians, statisticians, external sources)
- Complied with other plagiarism criteria specified in the Programme Handbook
- I understand that any false claim for this work will be penalised in accordance with the University regulations
- Received ethical approval from the School of Health in Social Science, University of Edinburgh

OR

- Received ethical approval from an approved external body and registered this application and confirmation of approval with the School of Health in Social Science's Ethical Committee

Signature

Date

Please note:

- If you need further guidance on plagiarism, you can:
 - Speak to your personal tutor or supervisor

- View university regulations at <http://www.ed.ac.uk/schools-departments/academic-services/policies-regulations>
- Referencing for most assessed work should be in the format of the BPS style guide, which is freely available from the BPS web site

Acknowledgements

I would first like to thank each person who agreed to take part in this study and offering up their time. I would like to also say a special thanks to all the health professionals and facilitators who helped with recruitment, without their support and hard work, this study would not have been possible.

Thank you to my supervisors Dr Paul Morris, Dr Nuno Ferreira, Dr Laura Keith and Line Manager Frances Baty for your guidance, encouragement and support throughout this process. I wish to thank Marie and Niamhy for co-rating papers and Tara Graham for generously sharing her wealth of knowledge on all things research! Thank you to my study buddy Patrick- every thesis needs one!

Finally, I would like to thank my family for always encouraging me to follow my passion regardless of any obstacles and ensuring that I always had an unlimited supply of Lyon's tea- each cup felt like a warm hug from home! To my friends both near and far for your patience and good humour and ability to remind me of things outside of thesis work. A special thank you to Andrew who has been a steadying force with his care, support and wise counsel and for reminding me that every journey has a destination and always keeping this one in sight for me- it's been invaluable.

List of Tables

Table 1.1 Electronic Database Search Terms

Table 1.2 Ratings of Study Quality for Included Studies

Table 1.3 Summary of studies included in review

Table 1.4 FM Participants Meta-analysed Pooled Mean Scores of the SF-36 subdomains (each subscale score from 0-100; higher scores indicates better HRQoL)

Table 1.5 Meta-analysed Differences between FM and HC groups on SF-36 subgroup scores

Table 2.1 Mean Differences between Genders on Outcome Measure Scores

Table 2.2 Pearson's Correlation Coefficients for Correlations of Key Variables, Age and Pain Duration

Table 2.3 Summary of Hierarchical Regression to Predict Quality of Life

Table 2.4 Summary of Hierarchical Regression to Predict Pain Disability

List of Figures

Figure 1.1 Search Process

Figure 1.2 Meta-analysed Differences between PCS and MCS for FM Individuals

Figure 1.3 Meta-analysed Difference between PCS and MCS in FM group compared to HC group

List of Appendices

Appendix A - Full Author Guidelines for 'Journal of Contextual Behavioural Science'

Appendix B - Quality rating tool for Systematic Review

Appendix C - Excluded studies from Systematic Review

Appendix D - Systematic review data extraction form

Appendix E - Forest Plots of Pooled Means for FM on subdomains of the SF-36

Appendix F- Forest Plots of Differences between Means for FM and HCs on SF-36 subdomains

Appendix G - Sensitivity and Subgroup Analysis Table

Appendix H - Full Author Guidelines for 'The Clinical Journal of Pain'

Appendix I- Permissions for Questionnaires

Appendix J – IRAS Protocol for Empirical Study

Appendix K – Ethical Approval Letters

Appendix L - Participant Information Sheet

Appendix M – Demographic Questionnaire

List of Abbreviations

FM- Fibromyalgia	FAS-Fibromyalgia Assessment Status
HC- Healthy Control	FIQ- Fibromyalgia Impact Questionnaire
HRQoL- Health Related Quality of Life	VAS- Visual Analogue Scale
SF-36- Short Form 36	QOL-Quality of Life
PF-Physical Functioning	
PR- Physical Role	
BP- Bodily Pain	
GH- General Health	
VT- Vitality	
ER- Emotional Role	
SF- Social Functioning	
MH- Mental Health	
CI- Confidence Interval	
MCS- Mental Component Summary	
PCS- Physical Component Summary	
BMI- Body Mass Index	
ACR- American College of Rheumatology	
SCS- Self-Compassion Scale	
BPI-SF- Brief Pain Inventory- Short Form	
RSES- Rosenberg Self-Esteem Scale	
HADS- Hospital Anxiety and Depression Scale	
PDQ- Physical Disability Questionnaire	

Word Counts:

Full Thesis Abstract- 592

Journal Article 1-6631

Journal Article 2-7858

Total Word Count-15077

Full Thesis Abstract

Background: Chronic pain conditions particularly fibromyalgia (FM) are an important issue considering the demand that they place on health services and the negative impact on an individual's well-being. Research regarding what impact FM may have on the various domains of health related quality of life is sparse. In addition, research identifying predictors relating to a person's self-concept on pain outcomes such as quality of life and disability is scarce in a chronic pain population. Self-esteem has been linked to psychological adjustment in various chronic diseases (Chong et al, 2009) and is suspected to be a vulnerability factor in FM (Johnson et al., 1997; Michielsen et al., 2006). Research on self-compassion found that it is a significant predictor of increased psychological adjustment in people with chronic medical conditions (Wren, Somers, & Wright, 2011). The majority of research on self-compassion has been conducted in non-clinical samples. Identifying the predictive values of self-esteem and the self-compassion subscales (self-warmth and self-coldness) in relation to pain outcomes are useful provided they can be enhanced and utilised to inform intervention.

Aims: This research portfolio had two aims, to systematically review and carry out a meta-analysis with the literature investigating the impact of FM on health related quality of life using the Short Form-36 version one and to investigate whether self-warmth, self-coldness and self-esteem act as predictors of quality of life and disability for individuals with chronic pain.

Method: A systematic review and meta-analysis of studies which assessed the impact of FM on health related quality of life was undertaken. Sensitivity and subgroup analyses were conducted to address the level of heterogeneity in the studies. The empirical study comprised of a cross-sectional design in which 60 individuals with chronic pain were recruited from three NHS sites based in Fife and Lothian and a pain self-management group in Fife to complete six validated psychometric questionnaires: Self Compassion Scale (Neff, 2003), Rosenberg Self-Esteem Scale (Rosenberg, 1965), Brief Pain Inventory-Short Form (Cleeland, 1991), Pain Disability Questionnaire (Anagnostis, Gatchel, & Mayer, 2004), Hospital Anxiety and Depression Scale (Zigmond and Snaith, 1983) and Quality of Life Scale (Burckhardt, Woods, Schultz, & Ziebarth, 2003).

Systematic Review Results: In total, 18 studies met the inclusion criteria of the systematic review and demonstrated that the health related quality of life of individuals with FM was lower compared to healthy controls on all eight subdomains of the SF-36 particularly Physical Role. It was found that Social Functioning was the subdomain least affected when comparing participants with FM to healthy controls.

Empirical Project Results: Quality of life was lower than previous research with females reporting a higher level of quality of life than males. It was found that after controlling for demographic and clinical variables, self-warmth was the only significant predictor for quality of life but not on levels of disability. Neither self-coldness nor self-esteem were significant predictors on scores of quality of life or disability.

Conclusion: The above studies expand literature on the nature of chronic pain and its outcomes such as health related quality of life. The meta-analysis evidenced that HRQoL was lower in individuals with FM than in healthy controls. The association between symptom severity and quality of life requires further investigation in FM.

Psychological interventions targeting the development of self-warmth attributes and skills may have a beneficial effect in improving quality of life for people with chronic pain. While the systematic review highlighted that FM has a more significant impact physically rather than mentally in relation to quality of life, both studies highlighted the need for interventions to target the psychological adjustment of people with chronic pain conditions.

Lay Summary of Thesis

This thesis consists of two papers. The first paper looks at the effect that fibromyalgia has on people's quality of life. Fibromyalgia is a long term condition that causes pain all over the body. Quality of life was measured by a questionnaire called the Short-Form-36. This questionnaire focuses on eight different areas of quality of life known as subdomains regarding ability to carrying out physical activities, general health, pain, mental well-being, ability to interact socially, and energy levels. The first paper looked at 18 studies which compared the scores of the SF-36 subdomains for people with fibromyalgia to people who do not have any pain conditions. People who do not have any pain condition are referred to as healthy controls in the present study. This study combined all scores of the 18 studies for the SF-36 subdomains together to get an average score in order to see which one would be most affected in people with fibromyalgia compared to healthy controls. It was found that people with fibromyalgia reported poorer quality of life than healthy controls particularly in relation to their physical ability to carry out activities. Ability to interact socially was least affected in people with fibromyalgia compared to healthy controls. Overall it was concluded that people with fibromyalgia are not all similar with regards to levels of quality of life and reasons for this should be explored further in future research. The second paper in this thesis focuses on people with chronic pain, which means that they have had continuous pain for at least three months. This paper aimed to answer whether an individual's sense of worth (self-esteem) or an ability to be kind to oneself (self-warmth) or judgemental to oneself (self-coldness) in times of difficulty predicted pain outcomes in relation to their levels of quality of life and level of disability. Sixty Individuals with chronic pain completed six questionnaires which asked questions relating to their severity of pain, mood, quality of life, disability, self-esteem, self-warmth and self-coldness. It was found after carrying out statistical analysis that people with higher levels of self-warmth regardless of mood or pain had better levels of quality of life but it did not affect levels of disability. However, self-coldness and self-esteem did not predict better or poorer levels of quality of life or disability. From the results of this study, it was suggested that psychological interventions should be aimed at helping to increase people's self-warmth in order to improve levels of quality of life.

Journal Article 1

The impact of fibromyalgia on health related quality of life using the Short-Form 36: A systematic review and meta-analysis

Authors

Caitríona Doughty, Dr. Laura Keith, Dr. Paul G. Morris, & Dr. Nuno Ferreira

Affiliations

Department of Clinical Psychology NHS Fife, Clinical and Health Psychology, University of Edinburgh Scotland.

Corresponding Author

Caitríona Doughty, Clinical Psychology Department, Lynebank Hospital, Halbeath Road, Dunfermline, Fife, KY11 4UW

Written in accordance with the instructions for 'Journal of Contextual Behavioural Science' (see Appendix A for author guidelines)

Abstract

Objective: To systematically review the literature to identify the impact of fibromyalgia (FM) on individuals with fibromyalgia in comparison to aged matched healthy controls on specific aspects of health related quality of life (HRQoL) using the Medical Outcomes Study Short-Form-36 (SF-36).

Methods: Multiple electronic databases were searched for studies reporting SF-36 subdomain mean and standard deviation scores for people with FM and healthy controls. Studies were methodologically appraised using a specifically developed quality criteria tool. A random-effects meta-analysis was performed to address the review questions.

Results: In total, 18 studies were eligible for inclusion in the meta-analysis, reporting on 1733 individuals with FM and 3468 healthy controls. Meta-analyses found that the subdomain Physical Role was the most severely effected in individuals with FM compared with Healthy Controls. The least affected subdomain was found to be Social Functioning (SF). It was calculated that the meta-analysed pooled mean HRQoL score for the SF-36 physical component summary score was 38.81 (95% CI: 31.2-46.4) and mental component summary score was 42.3 (95% CI: 34.3-50.2). Sensitivity analyses in relation to the methodology quality criteria did not significantly reduce heterogeneity in the review. Subgroup analysis did find a reduction in heterogeneity in relation to geographical area.

Conclusions: Individuals with FM have lower levels of HRQoL than healthy controls. This review supports that FM seems to have a greater negative impact on physical rather than on mental component summary scores of HRQoL though both are substantially affected.

1.1 Introduction

Fibromyalgia (FM) is a chronic pain disorder that affects 2-3% of the general population and is more prevalent in middle aged women (Quiroz, 2013). The prevalence of fibromyalgia can vary depending on different countries (Branco, Bannwart, Failde et al., 2010). Similarly, the particular characteristics and symptoms reported by individuals with fibromyalgia may differ depending on geographical regions which has been found among patients with chronic widespread musculoskeletal pain (McBeth & Jones, 2006). The American College of Rheumatology (ACR; 1990) developed criteria to classify FM for research, which included the presence of chronic widespread pain in combination with tenderness on examination at 11 or more 18 anatomical sites known as tender points. In 2010, the ACR introduced new preliminary diagnostic criteria which no longer requires tender point examination. Although FM is characterised by widespread pain, other symptoms include fatigue, sleep disturbance, morning stiffness, headaches and depressive mood (Wolfe, Clauw & Fitzcharles, 2010). The aetiology of FM is unknown, although it is generally considered a central sensitization syndrome (Clauw, 2014). This refers to the theory that the central nervous system may amplify sensory input across various organ systems resulting in a heightened sense of pain in a number of areas e.g. headaches, irritable bowels, tender points (Woolf, 2010). There is no known cure for FM, therefore, treatment is focused on preserving or improving daily function and alleviating pain (Albin, Fitzcharles, & Buskila, 2013).

FM has diverse effects on health-related quality of life (HRQoL), both physically and mentally (Hoffman & Dukes, 2008). Various terms have been used interchangeably with HRQoL including functional status, health status, and quality of life (Van Son, De Vries, Roukema, & Den Oudsten, 2013). An international group of researchers suggest that there are six fundamental domains of HRQOL: physical functioning, psychological functioning, social functioning, role activities, overall life satisfaction and perceptions of health status (Berzon, Hays, & Schumaker, 1993). While the objective dimensions aim to define a patient's degree of health, the individual's subjective evaluation serves to translate the health status into the actual quality of life experienced. Evaluating HRQOL

can be useful in that reduced HRQoL in FM is associated with increased use of healthcare resources (Kim, Kim, & Lee, 2012), poor prognosis of treatment (Torre, Martin-Corral, & Callejo, 2008) and increased levels of depression (Soriano-Maldonado, Amris, & Ortega, 2015). Therefore, decreasing the adverse impact of FM on HRQoL should be a key focus for informing treatment development.

The Medical Outcome Study 36-item Short Form Health Survey (SF-36) is one of the most widely used tools for assessing HRQoL (Garratt, Ruta, Abdalla, Buckingham & Russell, 1993). It defines HRQoL as the degree to which physical health affects an individual's perceived functioning and well-being in relation to psychological, social and physical aspects of life (Ware & Sherbourne, 1992). The SF-36 is organised into eight subscales that evaluates: physical functioning (PF), physical role (PR), bodily pain (BP), general health (GH), vitality (VT), emotional role (ER), social functioning (SF) and mental health (MH). Scores on these subscales can be combined to form two higher-order summary scores, the Physical Component Summary (PCS) and Mental Component Summary (MCS). The PCS is calculated by positively weighting the four subscales in the physical domain (PF, PR, BP and GH) and the remaining mental domain subscales (VT, SF, ER, MH) negatively. In contrast the MCS is calculated by positively weighting the four mental domain subscales and negatively weighting the four physical domain subscales. This measure has been found to be a reliable and a valid measure in FM (Timmerman, Calfa, & Stuifbergen, 2013), correlating well with disease-specific measures like the Fibromyalgia Impact Questionnaire-Revised (FIQ-R; Bennett, Friend, & Jones, 2009) and the Fibromyalgia Assessment Questionnaire (FAS; Salaffi, Sarzi-Puttini, & Girolimetti, 2009). Research suggests that the SF-36 is a useful generic measure which can compare HRQoL between populations rather than providing a disease specific assessment such as the FIQ and the FAS.

Increased levels of pain, physical disability, and mood disorders (anxiety and depression) have been associated with reduced quality of HRQoL for individuals with FM (Campos, 2012). Studies suggest that several factors may affect HRQoL in individuals with chronic pain, such as gender, age, body mass index (BMI), disease severity, and duration of disease (Moradkhani, Beckman, & Tabibian, 2013; Faust, Halpern, Danoff-Burg, & Cross, 2012). To date there have been a limited number of systematic reviews looking at the

impact of FM on HRQoL. Hoffman et al. (2008) has been the most recent systematic review which looked at health related quality of life in people with fibromyalgia compared to that of people in the general population and people with other long term conditions using the SF-36 and SF-12. While Hoffman's (2008) review found that individuals with FM had significant impairments in both mental and physical health domains in comparison to the general population and individuals with specific pain conditions, it did not focus on the eight components of the SF-36 or the two higher order summary scores. Therefore, the present study will be the first to carry out a meta-analysis examining which specific components of the SF-36 are most effected for individuals with FM. Such information can highlight the specific areas where individuals with FM experience particular difficulty and allow healthcare providers to develop interventions for these specific difficulties.

1.1.1 Aims of the Research:

1. Systematically review the literature appraising the impact of FM on the eight subdomains of the SF-36 by providing a pooled meta-analysed mean score for each subdomain of the SF-36.
2. To compare scores of individuals with FM on the SF-36 with aged matched healthy controls by providing a pooled meta-analysed mean difference score for each subdomain.
3. To compare both meta-analysed pooled mean scores of Physical Component Scores (PCS) and Mental Component Scores (MCS) for individuals with FM to determine which is more severely affected by FM.
4. To compare scores of individuals with FM on the PCS and MCS with aged matched healthy controls.

1.2 Materials and Methods

1.2.1 Search Strategy

In order to identify eligible studies, systematic searches of the following databases were conducted: Web of Science Core Collection, Ovid (incorporating EMBASE Classic and

Embase 1947 to 2015 October 07, Psych Info 1806 to October Week 1 2015, Ovid Medline In-Process and Other Non-Indexed Citations and Ovid Medline (R) 1946 to Present) and EBSCOHost (incorporating CINAHLPlus and ERIC) online databases. Search terms were developed (Table 1.1) with reference to common key words and were searched for within titles, keywords, abstracts. Since the SF-36 was developed in 1992, the search was restricted from January 1992 to October 2015. Following search completion, titles and abstracts were initially screened to eliminate studies not meeting the inclusion criteria. The remaining studies were read in full to determine appropriateness for inclusion.

Table 1.1 Electronic Database Search Terms

Electronic Database Search Term	
(1)	Fibromyalgia
(2)	Health Related Quality of Life
(3)	Quality of Life
(4)	Health Status
(5)	HRQoL
(6)	QoL
(7)	Short Form 36
(8)	SF-36
(9)	SF36
(10)	(2) or (3) or (4) or (5) or (6)
(11)	(1) and (10)
(12)	(7) or (8) or (9)
(13)	(11) and (12)

1.2.2 Selection Criteria

Studies were eligible for selection if they used quantitative methods and analysis, which focused on adult samples (defined as age > 18 years) with a diagnosis of FM based on either the 1990 or 2010 published classification criteria by the American College of Rheumatology (ACR). The review also required that the studies used the SF-36 version 1 to assess HRQoL rather than the SF-36 version 2 as they have different scoring systems which limits the comparability of the scores. While SF-36 version 2 has shown improvement over accuracy, reliability and validity compared to version 1, many studies in relation to rheumatic disorders and FM continue to use version 1 (Matcham, Scott, & Raynor, 2014; Wolfe, Michaud, Li, &

Katz, 2010). In order to reduce publication bias, grey material including academic, doctoral and masters' dissertations and other unpublished papers were included. Non empirical work such as book chapters or reviews were excluded. Addressing the review question involved including only case-controlled studies so that the HRQoL subdomain scores could be compared between individuals with FM and the general population. Theoretical or review studies were excluded.

1.2.3 Data Extraction and Quality Assessment

It is important to evaluate the methodological quality of studies in the systematic review so that the reliability and validity of the result can be predicted (Jarde, Losilla & Vives, 2013). Available tools for evaluating methodological quality are predominantly developed to evaluate treatment effectiveness using a randomised controlled trial design e.g. the Centre for Reviews and Dissemination (CRD, 2008). As the eligible studies in this review are observational studies, such checklists would not be compatible. While reporting on quality of observational studies has increased in recent times (Mallen, Peat, Croft, 2006) systematic reviews on quality assessment tools for evaluating observational studies have concluded that no single tool has been found to be reliable (Sanderson, Tatt, & Higgins, 2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE; Vandembroucke, von Elm, Altman, et al, 2007; vonElm, Altman, Egger et al, 2007) is supported by the Cochrane Collaboration and contains a checklist of recommendations of items to be included in reports of observational studies in epidemiology. However, it was not designed to evaluate methodological quality, therefore, for this review an assessment tool was developed and adapted from a number of sources including the STROBE (Kmet, Lee, & Cook, 2004), York's Centre for Reviews and Dissemination guidance for undertaking reviews in healthcare (CRD, 2008) and Scottish Intercollegiate Guidelines Network (SIGN, 2008). The final assessment tool (Appendix B) has a total of five quality criteria which include the domains of selection bias, statistical power, measure validity and method of analysis. Four categories were arbitrarily established for quality assessment: 'yes=2', 'partial=1', and 'no or not applicable=0'. In keeping with the guidance of the CRD (2008), a total numerical score was not calculated for the quality of each paper (Juni, Witschi, Bloch, & Egger, 1999). All studies were rated according to the quality criteria by the author and two independent raters. Reliability was good with a Krippendorff's alpha of .7 (Hayes & Krippendorff, 2007).

Discrepancies were resolved through discussion and re-examination of the studies. The quality ratings assigned for each paper are summarised in Table 1.2. In an attempt to retrieve missing or incomplete data particularly MCS and PCS scores, emails were sent to all authors of the studies that were included in this review.

1.2.4 Statistical Analysis

The eight subdomain scores of the SF-36 for individuals with FM and control groups in each study were extracted as mean totals and mean differences (MD) \pm SD. In addition, pooled means for PCS and MCS scores in the FM sample were meta-analysed for six of the studies that reported these values. Mean differences for the PCS and MCS were calculated and compared within the group of individuals with FM to determine whether there was a significant difference between the two components in this group. The PCS and MCS scores were also compared between individuals with FM and the healthy controls using mean totals and standard deviations. The OpenMeta-Analyst (Wallace, Dahabreh, Trikalinos et al., 2012) programme calculated scores in all studies eligible for the meta-analysis with 95% confidence intervals and combined by using random effects model (DerSimonian-Laird random effects method). Heterogeneity was assessed using the value of I^2 , whereby 25%, 50% and 75% indicated low, moderate and high heterogeneity respectively (Higgins, Thompson, Deeks, & Altman, 2003). Heterogeneity was found to be high between studies, therefore sensitivity and subgroup analyses were conducted.

1.3 Results

1.3.1 Search Results

Electronic databases were searched on 7th October 2015, with a return of 986 studies. Following removal of duplicates, there were 633 studies. First, titles of articles were screened resulting in the exclusion of 183 studies. Secondly, abstracts were reviewed and 341 were removed due to not meeting eligibility criteria for inclusion. Reference lists of all 109 studies that met the inclusion criteria were read in full. The 91 studies that were excluded at this stage are outlined in Appendix C. Eighteen studies were identified that met the inclusion criteria and were deemed appropriate to be part of the final methodological review and

analysis. This process is summarised in Figure 1.1 by a flowchart based on the PRISMA statement.

1.3.2 Characteristics of Included Studies

Table 1.3 provides a summary of descriptive characteristics and key findings of the 18 studies included in this systematic review. The included studies were all case-controlled cross-sectional studies published between 2000 and 2015. All studies included in the review used the 1990 American College of Rheumatology. Studies differed by country of origin and used translated versions of the SF-36 including Turkish (Akkaya et al., 2012; Alanoglu et al., 2005; Batmaz et al., 2013; Birtane et al., 2007; Tander et al., 2008; Turkyilmaz et al., 2012; Tuzun et al., 2004), Portuguese (Avila et al., 2014; Cardoso et al., 2011; Pagano et al., 2004), Spanish (Carillo et al., 2015; Segura-Jimenez et al., 2015), Italian (Salaffi et al., 2009), Hebrew (Neumann et al., 2000), Persian (Kolahi et al., 2010; Roshan et al., 2008) and Swedish (Andrell et al., 2014).

1.3.3 Methodological Quality Assessment

The methodological quality scores for each paper are provided in Table 1.2. According to the quality assessment tool, the majority of the studies included were of good methodological quality scoring highly across all criteria apart from the power/sample size. Ten studies were considered to be under-power (Alangolu et al., 2005, Avila et al., 2014, Batmaz et al., 2013, Birtane et al., 2007, Cardoso et al., 2011, Pagano et al., 2004, Smith et al., 2010, Tander et al., 2008, Turkyilmaz et al., 2012, & Tuzun et al., 2004).

1.3.4 Sample characteristics

There was marked variation in sample sizes with the number of individuals with FM ranging from 16 (Cardoso et al., 2011) to 459 (Segura-Jimenez et al., 2015). There was a total of 1733 individuals with FM included in this review. The control group in this review had a sample size ranging from 16 (Cardoso et al., 2011) to 1579 (Salaffi et al., 2009), comprising of 3468 in total. It was found that 15 studies recruited only females with FM, two studies included males with FM (Avila et al., 2014; Salaffi et al., 2009) and one study only included males with FM (Batmaz et al., 2013). The mean age of participants in the reviewed studies were 46 years for both groups, however seven studies only provided a total age score for both groups

(Akkaya et al., 2012; Alanoglu et al., 2005; Andrell et al., 2014; Avila et al., 2014; Batmaz et al., 2013; Birtane et al., 2007; Smith et al., 2010). Overall studies tended not to include individuals over the age of 65. Five studies specified age restrictions such as 35-60 (Cardoso et al., 2011; Pagano et al., 2004; Roshan et al., 2008, Segura-Jimenez et al., 2015., Smith et al., 2010), however only one of these studies (Roshan et al., 2008) explored age as a confounding variable in relation to HRQoL.

The majority of studies also had a trained health professional confirm the FM diagnosis rather than relying solely on self-report measures. Symptom severity of FM was measured using various psychometric measures in the reviewed studies. The Fibromyalgia Impact Questionnaire (FIQ) was administered in seven of the studies (Akkaya et al., 2012; Avila et al. 2014, Birtane et al. 2007, Carrillo et al. 2011, Pagano et al. 2004, Tander et al. 2008, and Turkyilmaz et al. 2012), the Fibromyalgia Impact Questionnaire-Revised in one study (Segura et al., 2015) whereas a Visual Analogue Scale, tender point count and less well defined yes/no checklist of chronic symptoms were used in the remaining studies. Additionally, given the variation in measures used, the specific symptoms investigated within each of the individual studies also varied as did the intensity of the symptoms presented.

1.3.5 Settings

Individuals with FM were recruited from outpatient clinics and FM associations. Two of the studies included in the review used general population-based samples rather than healthcare settings (Salaffi et al. 2009; Smith et al., 2010). A benefit of using a general population sample was that results were not biased by help-seeking (Mas, Carmona, Valverde, 2008).

1.3.6 Statistical Analysis

A variety of statistical analyses were conducted across studies and ten of the studies had normally distributed data. T-tests and analyses of variance were the prominent statistical techniques to examine the difference between individuals with FM and healthy controls in relation to the SF-36 sub-domains. Four studies were awarded “yes” and fourteen received “partial” in relation to appropriate use of statistics. The studies that were awarded “yes” reported confidence intervals and effect sizes. Reporting of demographic information was variable. Some studies noted significant differences between individuals with FM and healthy controls in relation to education level, BMI and employment which were not accounted for

in their analysis of SF-36 scores (Kolahi et al, 2010; Pagano et al. 2004). In order to explain the level of heterogeneity found, sensitivity analysis was conducted.

Figure 1.1 Search Process




Table 1.2 Ratings of Study Quality for Included Studies

	Sampling	Design & Method	Analysis		
	Recruitment Method	Fibromyalgia Measure	Power & Sample Size	Statistical Analysis	Confounding Variables
Akkaya, 2012	no	yes	partial	yes	yes
Alanoglu, 2005	no	no	no	partial	yes
Andrell, 2014	partial	no	yes	partial	no
Avila, 2014	no	yes	no	partial	partial
Batmaz, 2013	no	no	no	partial	no
Birtane, 2007	no	yes	no	partial	yes
Cardoso, 2011	no	no	no	yes	no
Carillo, 2015	no	yes	partial	partial	no
Kolahi, 2010	no	no	yes	no	yes
Neumann, 2000	no	no	yes	no	no
Pagano, 2004	no	yes	no	partial	partial
Roshan, 2008	no	no	yes	partial	yes
Salaffi, 2009	partial	no	yes	yes	yes
Segura, 2015	partial	yes	yes	yes	yes
Smith, 2010	partial	no	no	partial	yes
Tander,, 2008	no	yes	no	partial	yes
Turkyilmaz, 2012	no	yes	no	partial	yes
Tuzun, 2004	no	no	no	partial	yes

Table 1.3 Summary of studies included in review.

Author, Country	Year,	Sample Size	Recruitment	% female	Mean Age (SD)	Mean Disease Duration	Measure of FM	SF-36 PCS	SF-36 MCS
Akkaya et al., 2012 Turkey		FM= 51 CG= 41	Outpatient	FM=100 CG=100	Total Sample=35.5(9.9)	2.8(2.2)	FIQ FM= 70.8 (13.2) CG= 8.2 (9.6)		
Alanoglu et al., 2005 Turkey		FM= 34 CG= 22	Outpatient	FM=100 CG=100	Total Sample=37(10)	1.5(1)	TPC		
Andrell et al., 2014 Sweden		FM= 203 CG= 1055	Outpatient	FM=100 CG=52	Total Sample=46			FM=28.7(7.7) CG=50.0(9.4)	FM=38.0(13.7) CG=49.7(10.7)
Avila et al., 2014 Brazil		FM= 20 CG= 20	Outpatient	FM=95 CG=90	Total Sample=32.7(9.1)		FIQ FM=52(13.54) HC=26.0 (16.06)		
Batmaz et al., 2013 Turkey		FM= 37 CG= 35	Outpatient	FM=0 CG=0	Total Sample= 37.2(8.4)		VAS		
Birtane et al., 2007 Turkey		FM= 30 CG= 30	Outpatient	Not reported	Total Sample= 45.8(7.0)		FIQ		
Cardoso et al., 2011 Brazil		FM= 16 CG= 15	Outpatient	FM=100 CG=100	FM=53.5(7.5) CG=54.1(4.4)		SSB		
Carrillo et al., 2015 Spain		FM= 52 CG= 55	Outpatient	FM=100 CG=100	FM=46.2(9.4) CG=45.1(9.4)	6.5(5.0)	FIQ FM=62.9(16.2)		
Kolahi et al., 2010 Iran		FM= 100 CG= 100	Outpatient	FM=100 CG=100	FM=49.9(7.7) CG=46.6(6.5)		VAS	FM=60.5(9.4) CG=58.5(9.5)	FM=74.5(10.9) CG=69.8(11.9)
Neumann et al. 2000 Israel		FM= 90 CG= 50	Outpatient	FM=100 CG=100	FM=48(10) CG=55(13)	10(3)	VAS		
Pagano et al., 2004 Brazil		FM= 40 CG= 40	Outpatient	FM=100 CG=100	FM=49.5(6) CG=49.5(6)		FIQ		
Roshan et al., 2008 Iran		FM= 80 CG= 74	Outpatient	FM=100 CG=100	FM=42.9(9.1) CG=37.2(10.3)	10(3)			

Salaffi et al., 2009 Italy	FM= 380 CG= 1579	General Population	FM=92 CG=50	FM=52.1(10.8) CG=53.9(12.9)	10.5(9.7)	VAS	FM=38.6(6.9) CG=49.7(9.0)	FM=32.8(10.9) CG=45.6(8.4)
Segura-Jimenez, 2015 Spain	FM= 459 CG=214	FM Association	FM=100 CG=100	FM=52.2 (7.1) CG=51.3(7.0)		FIQ-R FM=64 (14.7) CG=22.4 (15.6)	FM=29.9(8.4) CG=46.9(10.5)	FM=35.9(12.6) CG=47.7(18.9)
Smith et al., 2010 USA	FM= 41 CG= 44	General Population	FM=100 CG=100	Total Sample= 48.4 (6.9)			FM=32.4(19.6) CG=89.4(8.9)	FM=39.6(19.7) CG=85.0(9.5)
Tander et al., 2008 Turkey	FM= 30 CG= 30	Outpatient	FM=100 CG=100	FM=40.0(1.1) CG=40.4(1.0)	5.23(0.9)	FIQ=51.70(2.26)		
Turkilmaz et al., 2012 Turkey	FM= 37 CG= 31	Outpatient	FM=100 CG=100	FM=39.2(6.5) CG=39.1(6.2)		FIQ FM=64.8(15) CG= 21 (8.5)	FM=42.6(17.0) CG=57.6(17.3)	FM=49.3(18.5) CG=60.8(17.1)
Tuzun et al., 2004 Turkey	FM= 33 CG= 33	Outpatient	FM=100 CG=100	FM=41.6(7.3) CG=38.9(5.4)	8.6(6.6)	VAS		

Note. Sample Size: FM=Fibromyalgia; CG=Control Group; **Measure:** FIQ= Fibromyalgia Impact Questionnaire; FIQ-R= Fibromyalgia Impact Questionnaire-Revised; VAS= Visual Analogue Scale;

1.3.7 Meta-analysis of the eight subdomains of the SF-36

The pooled mean scores on the eight SF-36 subscales for individuals with FM (Table 1.4), with 95% confidence intervals (CI) indicates that the least affected subdomain on the SF-36 is Social Functioning. The worst affected subdomain for individuals with FM appears to be Physical Role. The subdomains related to physical health apart from Physical Functioning (Physical Role, Bodily Pain, and General Health) showed lower mean scores than the subdomains related to mental health apart from Vitality (Emotion Role, Social Functioning and Mental Health).

Table 1.4 FM Participants Meta-analysed Pooled Mean Scores of the SF-36 subdomains (each subscale score from 0-100; higher scores indicates better HRQoL)

SF-36 Subgroups	Mean	Confidence Intervals 95%	Heterogeneity I²
Physical Functioning	49.89	45.38 , 54.40	96.58
Physical Role	28.37	21.37, 35.37	98.14
Body Pain	33.98	29.70, 38.27	97.65
General Health	39.33	34.67, 43.98	97.81
Vitality	37.09	32.58, 41.60	98.64
Social Function	52.88	48.06, 57.70	97.34
Emotion Role Function	42.68	36.63, 48.74	96.03
Mental Health	50.81	46.99, 54.62	96.14

Table 1.5 presents the pooled mean differences between individuals with FM and healthy controls. Forest plots of each domain meta-analysis can be found in Appendix E. On all eight subscales of the SF-36, there were significant differences found between the FM group and

healthy controls with the healthy controls reporting a significantly better level of HRQoL. Appendix F includes forest plots for the differences between individuals with FM and healthy controls with regards to the meta-analysed SF-36 subdomains. The greatest difference appeared to be in relation to the subdomains of Physical Role, Emotional Role, Bodily Pain and Physical Functioning.

Table 1.5 Meta-analysed differences between FM and HC groups on SF-36 subgroup scores

SF-36 Subgroups	Mean Difference	Confidence Intervals 95%	Heterogeneity I²
Physical Function	-33.92	-40.31 , -29.53	95.87
Physical Role	-53.39	-62.92, -43.87	97.92
Body Pain	-38.23	-43.97, -32.48	96.94
General Health	-31.11	-35.91, -26.31	95.45
Vitality	-29.41	-34.59, -24.24	96.40
Social Function	-24.35	-30.53, -18.17	96.83
Emotion Role Function	-37.48	-45.92, -29.03	96.33
Mental Health	-21.24	-24.82, -17.65	91.57

1.3.8 Meta-analysis of the two subcomponents of the SF-36 (PCS and MCS)

The pooled overall scores for PCS and MCS for six studies that reported these components was 38.8 (95% CI: 31.2-46.4) and 42.3 (95% CI: 34.3-50.2). Figure 1.2. It was found that there were no significant differences between the PCS and MCS scores for individuals with FM (MD= 3.31; 95% CI, -9.45, 2.84 with high heterogeneity among studies I²= 97.72).

Figure 1.2 Meta-analysed difference between PCS and MCS for FM individuals





Figure 1.3 presents the forest plots representing the pooled mean differences for PCS and MCS between individuals with FM and healthy controls. Six studies compared scores on the PCS in individuals with FM and healthy controls. Individuals with FM had significantly lower scores on the PCS compared with healthy controls (MD-22.07; 95% CI, -28.95, -15.20 with high heterogeneity among studies $I^2=98.58$). Six studies compared scores on the MCS in individuals with FM and controls. Individuals with FM had significantly lower scores on the MCS compared with healthy controls (MD-16.59; 95% CI, -21.38, -11.80 with high heterogeneity among studies $I^2=94.74$). It can be seen that there were greater differences between the individuals with FM and healthy control on the PCS compared to the MCS.

Figure 1.3 Meta-analysed difference between PCS and MCS in individuals with FM compared to healthy controls



1.3.9 Sensitivity and subgroup analysis

Appendix G presents the pooled mean differences between individuals with FM and healthy controls on the SF-36 sub-domain scores according to each sensitivity and subgroup analysis. Sensitivity analysis explored the exclusion of studies that did not receive a 'yes' or in some cases 'partial' on the methodology quality criteria in relation to sampling methods; measures of FM symptom severity, power/sample size; reporting effect sizes and p-values; and controlling for confounding variables. The subgroup analyses were conducted according to area of origin.

The sensitivity analysis found that the level of heterogeneity remained in the high range ($I^2 \geq 75$) despite excluding studies based on the quality of the methodology. After excluding 14 studies for using convenience sampling, the four studies that were awarded 'partial' (Andrell, 2014; Salaffi et al., 2009; Segura-Jimenez et al., 2015; Smith et al., 2010) showed a reduction in heterogeneity on BP ($I^2=11.1$) and MH ($I^2=0$) sub-domains. The subgroup analysis indicated that area of origin did reduce heterogeneity: on PF ($I^2=0$), PR ($I^2=0$), GH ($I^2=50.1$), VT ($I^2=0$), ER ($I^2=0$) for Iran/Israel (Kolahi et al., 2010; Roshan et al., 2008); on PF ($I^2=49.39$), BP ($I^2=43.25$), and MH ($I^2=49.34$) for America/South America (Avila et al., 2014; Cardoso et al., 2011; Pagano et al., 2004; Smith et al., 2010); and on MH ($I^2=0$) for Europe/Scandinavia (Andrell et al., 2014; Carrillo et al., 2015; Salaffi et al., 2009; Segura-Jimenez et al., 2015). Heterogeneity remained in the high range for all Turkish (other) studies (Akkaya et al., 2012; Alanoglu et al., 2005; Batmaz et al., 2013; Birtane et al., 2007; Neumann et al., 2000; Tander et al., 2008; Turkyilmaz et al., 2012; Tuzun et al., 2004). Sensitivity and subgroup analysis were not conducted to explore heterogeneity for the PCS and MCS scores as there were only six studies (Egger, Smith, Schneider, & Minder, 1997).

1.3.10 Narrative synthesis of key findings

In total, 11 out of 18 studies included in the review reported a significant difference in all subdomains of the SF-36 between individuals with and healthy controls (Alanoglu et al., 2005; Andrell et al. 2014; Avila et al. 2014, Batmaz et al., 2013; Birtane et al., 2007; Carillo et al 2014; Pagano et al. 2004; Salaffi et al. 2009; Smith et al., 2010; Segura-Jimenez et al., 2015; Tuzun et al. 2004).

This meta-analysis calculated that Physical Role compared to the other eight SF-36 subdomains showed the greatest difference between individuals with FM and healthy controls. The forest plots (Appendix F) demonstrated that Pagano et al. (2004) compared to the other studies in the review reported the greatest difference between the FM group and the HC group in relation to PR. Pagano et al. (2004) matched controls for age and BMI, which received a 'partial' quality criteria rating for addressing confounding variables. The FIQ was also used but Pagano et al. (2004) did not report the total score of the FIQ or the mean differences between the FM and HC group to show the difference in severity of symptoms between the two groups. Turkyilmaz et al. (2012) on the other hand did not find any significant difference between the individuals with FM and healthy controls on the PR subdomain, however this study controlled for a number of variables which are known to impact quality of life e.g. BMI, marriage, education, occupation, economic status, mood. In term of addressing confounding variables on the methodology quality criteria, this study was awarded 'yes'. The FIQ was also used in this study and it was found that while there was a significant difference between individuals with FM and healthy controls, the total figures reported for the healthy controls were slightly higher than what has been previously reported in other studies (Akkaya et al., 2012; Sivas, Baskan, Aktekin, 2009). This may indicate that there was a slightly higher level of symptom severity in the control sample which may explain the lack of significant difference between individuals with FM and healthy controls in relation to this subdomain. This is in line with the finding that there is a strong correlation between the FIQ and the PR subdomain ($r=0.481$; $p<0.007$; Birtane et al., 2007). Cardoso et al., (2011) also did not report a significant difference on PR along with four other subdomains (VT, SF, ER, and MH). It must be noted that Cardoso et al., 2011 reported the smallest sample size ($N=16$) of all the studies in this review and received a 'no' rating on the methodology quality criteria for power. It is likely that the lack of significant differences between individuals with FM and healthy controls were a result of the study being insufficiently powered.

While this meta-analysis found PR to be the subdomain which was most negatively affected in the FM group compared to the HC group, the SF subdomain was reported to be the least negatively impacted between these two groups. Four of the 18 studies (Akkaya et al. 2012; Cardoso et al., 2011; Roshan et al., 2008; Turkyilmaz et al., 2012) reported that SF was not statistically significant between individuals with FM and healthy controls. Of particular

interest was Roshan et al., (2008) study whereby SF score was found to be higher in individuals with FM than the healthy controls. Roshan et al. (2008) controlled for marital status, education, occupation as well as age which may have resulted in them finding a similar SF and MH score in both individuals with FM and healthy controls. Two other studies in this review also found that MH was not statistically significant (Tander et al., 2008; Cardoso et al., 2011). Tander et al., (2008) assessed for depression by using the Beck Depression Inventory (BDI) whereby a score of >17 was considered to be a sign of depression. It may be that individuals with FM and healthy controls did not differ significantly on MH because the BDI eliminated the possibility of individuals with co-morbid depression.

Ten studies (Alanoglu et al., 2005; Andrell et al., 2014; Avilia et al., 2014, Birtane et al., 2007, Cardoso et al., 2011, Kolahi et al., 2010, Pagano et al., 2004, Roshan et al., 2008, Salaffi et al., 2009, Smith et al., 2010) in this review did not use any standardised depression measure to account for co-morbid depression which may increase the likelihood of a significant difference being present between individuals with FM and healthy controls on the MH subdomain as previous studies have indicated the high prevalence of co-morbid depression in individuals with FM (Fuller-Thomson, Nimignou-Young, Brennenstuhl, 2012). Two studies reported that when comparing individuals with FM and healthy controls they did not find a significant difference in all of the SF-36 domains, however these studies did not include p-values to specify which subdomains were insignificant (Kolahi et al., 2010; Neumann et al. 2000). For this reason, these two studies were awarded 'no' on the methodology quality criteria for statistical analysis as the lack of p-values limited the possibility of interpretation and comparability of results.

In terms of the PCS and MCS component scores it was found there were no significant difference between these components in individuals with FM. All five studies showed a slightly higher mean score for PCS compared to MCS except for Salaffi et al. (2009). It may be worth noting that apart from Segura et al. (2015), Salaffi seemed to receive the best ratings across all the methodology quality criteria. In their study they found that the PCS was influenced by widespread pain, education level and BMI whereas MCS was correlated with widespread pain, sleep abnormalities, physical functioning, fatigue, gender and low educational level. As Salaffi et al. (2009) used a general population, this sample may be more representative of individuals with FM as it reduces bias from help seeking (Mas et al., 2008).

Overall, the six studies seemed to confirm that the PCS is more severely affected in individuals with FM compared to healthy controls.

1.4 Discussion

Studies conducted across different geographical areas showed that individuals with FM had a consistent pattern of HRQoL impairment. Overall individuals with FM scored significantly lower on all eight health status domains compared with individuals in the general population. The results of this review show that individuals with FM have lower levels of HRQoL, in particular Physical Role. This is in line with previous literature which suggests that individuals with FM have poor scores on the physical dimensions of HRQoL (Campos et al., 2012). Social Functioning and Mental Health were the least affected subdomains in this review for individuals with FM. A study carried out by Alonso et al. (2004) that used the SF-36 across eight different countries found Social Functioning and Mental Health to be the least affected in relation to chronic illnesses. These results were consistent with a similar review carried out among individuals with rheumatoid arthritis (Matcham et al., 2014). The relatively low impact of these diseases on social functioning and mental health may be explained by individuals being able to adapt to the conditions of the disease and/or adopting new lifestyle behaviours. It may also be related to cultural aspects as it was found that social functioning was least affected in an Iranian study (Roshan et al., 2008). This is in line with other systematic reviews that have found social functioning to be the least affected subdomain in an Iranian population with rheumatoid arthritis (Shokri, Mottaghi, Qolipour, 2015) and an elderly population where it was suggested that individuals may be able to access social support from family and society due to the values and emphasis on community (Tajvar, Arab, Montazeri, 2008). One reason that Roshan may have found higher rates of SF than the control group was that the study controlled for marriage, education and occupation which have been found to be predictors of better related quality of life for Iranians with rheumatoid arthritis (Shokri et al, 2015). The greatest difference between individuals with FM and healthy controls were in the domains of Physical Role, Emotional Role, Bodily Pain and Physical Functioning. Other studies have suggested that the functional loss through incapacity to work and drop in income because of pain and fatigue may impact on a person's perceived health related quality of life (White, Speechley, Harth, & Ostbye, 1999).

The quality of studies appeared to be good, however the majority of studies used sampling criteria that limited their external validity due to a small number of studies investigating population based samples (Salaffi et al. 2009; Smith et al., 2010). Studies that used convenience sampling recruited participants from hospital outpatient clinics, which may represent the most affected individuals with FM. Therefore, these results may not be an accurate representation of the FM population (Mas et al., 2008). On the other hand, smaller sample sizes (Avila et al., 2014; Cardoso et al., 2011) may represent only patients who present themselves for research participation, who in turn, would usually represent a healthier patient group (Knudsen, Hotopf, Skogen, Overland, & Mykletun, 2010).

This review focused on one specific factor, HRQoL, however the majority of studies in this review explored a broad range of factors relating to FM such as auditory event related brain potentials (ERBP; Alangolu et al., 2005). While ERBP was relevant to FM, it was not related to HRQoL in FM. Therefore, this approach may not be the most appropriate as it may result in participant fatigue, with participants completing large batteries of questionnaires (Johnson et al, 2011). Moreover, it possibly will result in Type 1 error by increasing the likelihood of false positives. A small number of studies (Neumann et al., 2000) controlled for this by using adjustments to the criteria for significance such as the Bonferroni correction. However, this could increase the likelihood of Type II error (Perneger, 1998), a preferable alternative would be to conduct studies that included a smaller number of factors which may be more relevant to HRQoL.

Increased BMI, lower education levels, marriage status, lower employment have been associated with poorer SF-36 scores (Cherapanov, Palta, Fryback et al., 2011). These factors were controlled for in a number of studies, however only one study controlled for them using a multiple regression. For this reason, it was not possible to look at whether these factors predicted poor HRQoL in individuals with FM. Although individuals with FM and the control group were matched for age, some studies applied an age-criteria in order to match the groups (Cardoso et al., 2011; Pagano et al., 2004; Roshan et al., 2008; Segura-Jimenez et al., 2015), which may have introduced bias to the review. Overall studies tended not to include

participants over the age of 65, limiting the generalisability of the impact of FM on quality of life in older adults. This further reduces the interpretation and comparability of studies.

All studies used the ACR 1990 criteria for identifying FM rather than the 2010 preliminary criteria. While there is good agreement between the 1990 and 2010 criteria, it has been argued that the 2010 criteria have lower specificity and higher sensitivity (Goldenberg, 2009). This finding suggests that there is a higher possibility that individuals with other functional diseases but not generalised pain may be identified as having FM when using the 2010 ACR criteria (Carrillo et al., 2015). All of the studies in this review included predominantly female populations, which is in line with the literature reporting that 90% of individuals with FM are women (Yunus, 2002).

The high level of heterogeneity identified in the meta-analysis may be explained by variation in the study population including disease duration and symptom severity. While a number of studies accounted for variables such as gender, ethnicity, and marital status, a number of studies did not assess severity of symptoms. Eight studies in the review used a standardised measure for symptom severity and only six of these reported the values. Individuals with Fibromyalgia are a heterogeneous group and the rating number and impact of different symptoms are expected to vary between different populations (Branco et al, 2010). Research in this area has proposed a model of clustering individuals with FM into two categories based on physical symptoms and psychological symptoms of the FIQ (de Souza, Goffaux, Julien et al., 2009; Calandre, Garcia- Carrillo, Garcia-Leiva, 2011). Therefore, it would be useful to carry out a meta-analysis in the future looking at the impact of symptoms using the FIQ on the eight subdomains of the SF-36 to see if this may help to reduce the level of heterogeneity in the research.

Sensitivity analyses did not reveal that the results were sensitive to the study methodology except in relation to sampling. Random sampling or recruiting from a number of areas rather than convenience sampling seemed to reduce heterogeneity in BP and MH. It is surprising that heterogeneity was reduced in BP as pain tends to be variable among individuals with FM (Schaeffer, 1997) with average daily pain ratings varying in individuals depending on time of day (Okifuji, Bradshaw, Donaldson, 2011). Subgroup analysis in relation to geographical origin succeeded in reducing some of the heterogeneity. Reasons for this may be in relation to culture or the different translations of the SF-36; mean scores may not be comparable due to

different cultural variation in both phrasing of items and responses and in interpretation of the concept of HRQoL (Bullinger, Alonso, Apolone et al., 1998).

1.41 Strengths, limitation and implications for future research

Limitations of this review were that all studies were cross-sectional and therefore causation of poor HRQoL in individuals with FM could not be inferred. However, this review indicates that the physical role is most negatively affected in individuals with FM and provides a rationale upon which more methodologically rigorous experimental studies can be developed. In particular, it would be useful to look at what factors may be predictive of poor scores on physical role e.g. physical activity (Kop, Lyden, Berlin, Ambrose, Olsen, 2005) in individuals with FM which could help to develop intervention studies.

This study only considered studies that used the SF-36 to measure HRQoL, which may have excluded studies that were potentially relevant. For example, Mas (2008) recruited from the general population and compared HRQoL using the SF-12 in individuals with FM and healthy controls. This data would have been useful as the external validity of the study was strong, however, it would not have been possible to include in the meta-analysis.

A strength of the current review was that authors were contacted for further information such as PCS and MCS scores. Therefore, by providing the two higher summary scores along with subdomain scores, this review was able to show that the impact is significant in both physical and mental domains. The findings from this review show that while scores on the physical domains were poorer than the mental domains, treatment of FM should incorporate interventions that target helping people to improve both physical and mental components. This suggestion is in line with the SIGN Guidelines (2013) for chronic pain stipulating that a multidisciplinary team should be involved with chronic pain conditions. They also suggest that people with pain conditions tend to focus more on physical symptoms, therefore when referring patients for psychological assessment it should be clarified that the aim of a psychological intervention is to increase coping skills and improve quality of life.

Finally, many of the studies did not include information about severity of FM symptoms or mood ratings such as depression or anxiety, it was not possible to specify the association between the HRQoL and potential clinical factors contributing to low HRQoL. All of the above

limitations suggested, indicates that further research is needed to confirm the findings of this study.

In conclusion the review demonstrated that individuals with FM have a lower level of HRQoL particularly on the Physical Role domain. Considering the level of heterogeneity of the evidence, this conclusion should be interpreted with caution. As it was not possible to obtain individual patient data for this review, aggregate data was used, however individual patient data would be recommended for future research in this area as it can improve the quality of the data and the type of analyses that can be performed (Stewart & Tierney, 2002).

1.5 References

- Akkaya, N., Akkaya, S., Atalay., N., Balci, C., Sahin, F. (2012). Relationship between the body image and level of pain, functional status, severity of depression, and quality of life in patients with fibromyalgia syndrome. *Clinical Rheumatology*, 31, 983-988.
- Alanoglu, E., Ulas, U., Ozdag, F., Odabasi, Z., Cakci, A. (2005). Auditory event-related brain potential in fibromyalgia syndrome. *Rheumatology International*, 25, 345-349.
- Ablin, J., Fitzcharles, M.-A., Buskila, D., Shir, Y., Sommer, C., & Häuser, W. (2013). Treatment of Fibromyalgia Syndrome: Recommendations of Recent Evidence-Based Interdisciplinary Guidelines with Special Emphasis on Complementary and Alternative Therapies. Evidence-Based Complementary and Alternative Medicine, 2013
<http://doi.org/10.1155/2013/485272>
- Alonso, J., Ferrer, M., Gandek, B., Ware., J, Aaronson, N.K., Mosconi, P. et al. (2004). Health related quality of life associated with chronic conditions in eight countries: Results from the International Quality Life Assessment (IQOLA) Project. *Quality of Life Research*, 13, 283-29.
- Andrell, P., Schultz, T., Mannerkorpi, K., Nordeman, L., Borjesson, M., & Mannheimer, C. (2014). Health-related quality of life in fibromyalgia and refractory angina pectoris: A comparison between two chronic non-malignant pain disorders. *Journal of Rehabilitation Medicine*, 46, 341-347.
- Avila, L., de Araujo Filho, G., Guimaraes, E., Goncalves, L., Paschoalin, P (2014). Characterisation of the pain, sleep and alexithymia patterns of patients with fibromyalgia treated in a Brazilian tertiary centre. *Revista Brasileira de Reumatologia*, 54, 409-413.
- Batmaz, I., Sariyildiz, M., Dilek, B., Inanir, A., Demircan, Z. (2013). Sexuality of men with fibromyalgia: What are the factors that cause sexual dysfunction? *Rheumatology International*, 33, 1265-1270.
- Bennett, R.M., Friend, R., & Jones K.D. (2009). The revised fibromyalgia impact questionnaire (FIQR): Validation and psychometric properties. *Arthritis Research Therapy*, 11, R120

- Berzon, R., Hays, R.D., & Shumaker, S.A. (1993). International use, application, and performance of health-related quality of life instruments. *Quality of Life Research*, 2, 367-368.
- Birtane, M., Uzunca, K., Tastekin, N. Tuna., H. (2007). The evaluation of quality of life in fibromyalgia syndrome: A comparison with rheumatoid arthritis by using SF-36 Health Survey. *Clinical Rheumatology*, 26, 679-684.
- Branco, J.C. Bannwarth, B., Failde, I et al. (2010). Prevalence of fibromyalgia: Survey in five countries. *Seminars in Arthritis and Rheumatology*, 39, 448-453.
- Bullinger, M., Alonso, J., Apolone, G., Lepiege, A., Sullivan, M. (1998). Translating health status questionnaires and evaluating their quality: the IQOLA project approach. *International Journal of Quality of Life Assessment. Journal of Clinical Epidemiology*, 51, 913-923.
- Calandre, E.P., Garcia-Carrillo, J., Garcia-Leiva, J.M., Rico-Villademoros, F., Molina-Barea, R., et al. (2011). Subgrouping patients with fibromyalgia according to the results of the fibromyalgia impact questionnaire: a replication study. *Rheumatology International*, 31, 1555-1559.
- Campos, R.P., & Vasquez, M. (2012). Health-related quality of life in women with fibromyalgia: Clinical and psychological factors associated. *Clinical Rheumatology*, 31, 347-355.
- Cardoso, F., Curtolo, M. Natour, J., Junior, I. (2011). Assessment of quality of life, muscle strength and functional capacity in women with fibromyalgia. *Revista Brasileira de Reumatologia*, 51, 338-350.
- Carrillo-de-la-Pena, M.T., Trinanès, Y., González-Villar (2015). Convergence between the 1990 and 2010 ACR diagnostic criteria and validation of the Spanish version of the fibromyalgia survey questionnaire (FSQ). *Rheumatology International*, 35, 141-151.
- Centre for Reviews and Dissemination. (2008). Systematic Reviews: CRD's guidance for undertaking reviews in health care. CRD, University of York. York Publishing Services Ltd. DOI: https://www.york.ac.uk/media/crd/Systematic_Reviews.pdf
- Cherapanov, D. Palta, M., Fryback, D. Robert, S., Hays, R. & Kaplan, R. (2011). Gender differences in multiple underlying dimensions of health-related quality of life are associated with sociodemographic and socioeconomic status. *Medical Care*, 49, 1021-1030.

- Clauw, D. (2014). Fibromyalgia: A clinical review. *The Journal of the American Medical Association*, 15, 1547-1555.
- De Souza, J.B., Goffaux, P., Julien, N., Potvin, S., Charest, J., et al. (2009). Fibromyalgia subgroups: profiling distinct subgroups using the Fibromyalgia Impact Questionnaire. A preliminary study. *Rheumatology International*, 29, 509-515.
- DerSimonian, R., Laird, N. (1986) Meta-analysis in clinical trials. *Control Clinical Trials*, 7, 177-188.
- Egger, M., Smith, G.D., Schneider, M., Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315, 629-634.
- Faust, A., Halpern, L., Danoff-Burg, S., & Cross, R. (2012). Psychosocial factors contributing to inflammatory bowel disease activity and health-related quality of life. *Gastroenterology & Hepatology*, 8, 173-181.
- Fuller-Thomson, E., Nimigon-Young, J. & Brennenstuhl, S. (2012). Individuals with fibromyalgia and depression: Findings from a nationally representative Canadian survey. *Rheumatology International*, 32, 853-862.
- Garratt, A.M., Ruta, D.A., Abdalla, M.I., Buckingham, J.K., Russell, I.T. (1993). The SF-36 health survey questionnaire: An outcome measure suitable for routine use within the NHS? *British Medical Journal*, 306, 1440-1444.
- Goldenberg, DL (2009). Diagnosis and differential diagnosis of fibromyalgia. *American Journal of Medicine*, 122, S14-S21.
- Hayes, A.F. & Krippendorff, K. (2007). Answering the call for a standard reliability measure for coding data. *Communication Methods and Measure*, 1, 77-89.
- Higgins, J.P.T., Thompson, S.G., Deeks, J.J. & Altman, D.G. (2003). Measuring inconsistency in meta-analyses. *British Medical Journal*, 327, 557-560.
- Hoffman, D.L. & Dukes, E.M. (2008). The health status burden of people with fibromyalgia: a review of studies that assessed health status with the SF-36 or the SF-12. *International Journal of Clinical Practice*, 62, 115-126.

- Jarde, A., Losilla, J., Vives, J., & Rodrigo, M. (2013). Q-Cohh: A tool to screen the methodological quality of cohort studies in systematic reviews and meta-analyses. *International Journal of Clinical and Health Psychology*, 13, 138-146.
- Juni, P., Witschi, A., Bloch, R., & Egger, M. (1999). The hazards of scoring the quality of clinical trials for meta-analysis. *The Journal of the American Medical Association*, 282, 1054-1060.
- Kim, S., Kim, S., Lee, C.K., Lee, H.S. & Lee, S.H. (2012). Effect of fibromyalgia syndrome on the health-related quality of life and economic burden in Korea. *Rheumatology*, 52, 311-320.
- Kmet, L.M., Lee, R.C., & Cook, L.S. (2004). Standard quality assessment criteria for evaluating primary research papers from a variety of fields. Edmonton: Alberta Heritage Foundation for Medical Research (AHFMR). HTA Initiative #13. 2004. DOI: <http://www.ihe.ca/documents/HTA-FR14.pdf>
- Knudsen, A.K., Hotopf, M., Skogen, J.C., Overland, S., Mykletun, A. (2010). The health status of non-participants in a population based health study: The Hordaland health study. *American Journal of Epidemiology*, 172, 1306-1314.
- Kolahi, S., Khbazi, A., Hajaliloo, M., Namvar, L., Farzin, H. (2010). The evaluation of quality of life in women with rheumatoid arthritis, osteoarthritis and fibromyalgia as compared with quality of life in normal women. *The Internet Journal of Rheumatology*, 1, 1-5.
- Kop, W. J., Lyden, A., Berlin, A. A., Ambrose, K., Olsen, C., Gracely, R. H., Williams, D. A. and Clauw, D. J. (2005), Ambulatory monitoring of physical activity and symptoms in fibromyalgia and chronic fatigue syndrome. *Arthritis & Rheumatism*, 52: 296-303
- Mallen, C. Peat, G., Croft, P. (2006) Quality assessment of observational studies is not commonplace in systematic reviews. *Journal of Clinical Epidemiology*, 59, 765-769.
- Mas, A.J., Carmona, L. Valverde (2008). Prevalence and impact of fibromyalgia on function and quality of life in individuals from the general population: Result from a nationwide study in Spain. *Clinical and Experimental Rheumatology*, 26, 19-526.
- Matcham, F., Rayner, L., Hutton, J., Monk, A., Steel, C., Hotopf, M. (2014). Self-help interventions for symptoms of depression, anxiety and psychological distress in patients with physical illnesses: A systematic review and meta-analysis. *Clinical Psychology Review*, 34, 141-157.

- McBeth, J. Jones, K. (2006). Epidemiology of chronic musculoskeletal pain. *Best Practice Research of Clinical Rheumatology*, 21, 403-425.
- Moradkhani, A., Beckman, L.J. & Tabibian, J.H. (2013). Health-related quality of life in inflammatory bowel disease: Psychosocial, clinical, socioeconomic, and demographic predictors. *Journal of Crohn's and Colitis*, 7, 467-473.
- Neumann, L., Berzak, A., & Buskila, D. (2000). Measuring health status in Israeli patients with fibromyalgia syndrome and widespread pain and healthy individuals: Utility of the Short-Form 36-item health survey (SF-36). *Seminars in Arthritis Rheumatism*, 29, 400-408.
- Okifuji, A. Bradshaw, D.H., Donaldson, G.W. (2011). Sequential analyses of daily symptoms in women with fibromyalgia syndrome. *Journal of Pain*, 12, 84-93.
- Pagano, T., Matsutani, L., Ferreira, E., Marques, A., Pereira, C. (2004). Assessment of anxiety and quality of life in fibromyalgia patients. *Sao Paulo Medical Journal*, 122, 252-258.
- Perneger, T.V. (1998). What's wrong with Bonferroni adjustments? *The British Medical Journal*, 316, 1236-1238.
- Quiroz, L.P. (2013). Worldwide epidemiology of fibromyalgia. *Current Pain and Headache Reports*, 17, 356-342.
- Roshan, R., Shariatpanahi, S., Tavoli, A., Qafari, Z., & Fathi-Ashtiani, A. (2008). *Quality of life in women with fibromyalgia. Archives of Medical Science*, 5, 267-270.
- Salaffi, F., Sarzi-Puttini, P., & Girolimetti, R. (2009). Development and validation of the self-administered Fibromyalgia Assessment Stats: A disease specific composite measure for evaluating treatment effect. *Arthritis Research & Therapy*, 11, R125.
- Sanderson, S., Tatt, I.D., Higgins, J.P. (2007). Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: A systematic review and annotated bibliography. *International Journal of Epidemiology*, 36, 677-678.
- Schaefer, K.M. (1997). Health patterns of women with fibromyalgia. *Journal of Advanced Nursing*, 26, 565-571.
- Scottish Intercollegiate Guidelines Network (SIGN). SIGN 50 Guideline Developer's Handbook 2008. DOI: <http://www.sign.ac.uk/guidelines/fulltext/50/section1.html>

Scottish Intercollegiate Guidelines Network (SIGN). Management of Chronic Pain. Edinburgh: SIGN; 2013. (SIGN publication no. 136). Available from [url:http://www.sign.ac.uk/guidelines/fulltext/136/index.html](http://www.sign.ac.uk/guidelines/fulltext/136/index.html)

Segura-Jimenez, V., Alvarez-Gallardo, I., Carbonell-Baeza, A., Aparicio, V.A., & Ortega, F. et al. (2015). Fibromyalgia has a larger impact on physical health than on psychological health yet both are markedly affected: The al-Andalus project. *Seminars in Arthritis and Rheumatism*, 44, 563-570.

Shokri, A, Mottaghi, P., Qolipour, K. (2015). Quality of life and its predictors among Iranian patients with rheumatoid arthritis: A systematic review. *Jetnashapier Journal of Health Research*, 6, DOI: 10.5812/jjhr.6(2)2015.24636.

Sivas, F.A., Baskan, B.M., Aktekin, L.A. (2009). Depression in fibromyalgia patients' assessment of sleep disorders and quality of life. *Turkish Journal Physical Medical Rehabilitation*, 55, 8-12.

Smith, B.W., Papp, Z., Tooley, E.M., Montague, E., Robinson, A. (2010). Traumatic evets, perceived stress and health in women with fibromyalgia and healthy controls. *Stress and Health*, 26, 83-93

Soriano-Maldonado, A. Amris, K. & Ortega, F.B. (2015). Association of different levels of depressive symptoms with symptomatology, overall disease severity, and quality of life in women with fibromyalgia. *Quality of Life Research*, 24, 2951-2957.

Stewart, L. A., & Tierney, J. F. (2002). To IPD or not to IPD? Advantages and disadvantages of systematic reviews using individual patient data. *Evaluation and the Health Professions*, 25, 76-97.

Tajvar, M., Arab, M., Montazeri, A. (2008). Determinants of health related quality of life in elderly in Tehran, Iran. *BMC Public Health*, 8, 323-331.

Tander, B., Cengiz, K., Alayli, G., Ilhanli, I., & Canbaz, S. (2008). A comparison evaluation of health related quality of life and depression in patients with fibromyalgia syndrome and rheumatoid arthritis. *Rheumatology International*, 28, 859-865.

- Timmerman, G.M., Calfa, N.A. & Stufbergen, A.K. (2013). Correlates of body mass index in women with fibromyalgia. *Orthopaedic Nursing*, 32, 113-119.
- Torre, F. Martin-Corral, J., & Callejo, A. (2008). Health related quality of life and coping strategies with regard to pain in patients attended in a unit of pain. *Revista de la Sociedad Española del Dolor*, 15, 83-93.
- Turkyilmaz, A., Kurt, E., Karkucak, M., Capkin, E. (2012). Sociodemographic characteristics, clinical signs and quality of life in patients with fibromyalgia. *The Eurasian Journal of Medicine*, 44, 89-93.
- Tuzun, E., Albayrak, G., Eker, L., Sozays, S., Daskapan, A. (2004). A comparison study of quality of life in women with fibromyalgia and myofascial pain syndrome. *Disability and Rehabilitation*, 26, 198-202.
- Vandenbroucke, J.P., von Elm, E., Altman, D.G., Gøtzsche, P.D., Mulrow, C.D. et al. (2007). Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and Elaboration. *Annals of Internal Medicine*, 147, 163-194.
- Van Son, M.A.C., De Vries, J., Roukema, J.A. & Den Oudsten, B.L. (2013). Health status, health-related quality of life, and quality of life following ankle fractures: A systematic review. *International Journal of Care Injured*, 44, 1391-1402. Berzon
- Von Elm, E., Altman, D.G., Egger, M., Pocock, S.J., Gøtzsche, P.D., et al. (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Epidemiology*, 18, 800-804.
- Wallace, B., Dahabreh, I, Trikalinos, T., Lau, J., Trow, P., Schmid, C. (2012). Closing the gap between methodologists and end-users: R as computation back-end. *Journal of Statistical Software* 49, 1-15.
- Ware, J.E. & Sherbourne, C.D. (1992). The MOS 36-item Short-Form Health Survey (SF-36): I. Conceptual framework and item selection. *Medical Care*, 30, 473-83.
- White, KP., Speechley, M., Harth, M., Ostbye, T. (1999) The London Fibromyalgia Epidemiology Study: The prevalence of fibromyalgia syndrome in London, Ontario. *Journal of Rheumatology*, 26, 1570-1576.

- Wolfe, F., Clauw, D.J., Fitzcharles, M.A., Goldenberg, R.S., Katz, P. (2010). The American College of Rheumatology preliminary diagnostic critieria for fibromyalgia and measurement of symptom severity. *Arthritis Care and Research*, 62, 600-610.
- Wolfe, F., Michaud, K., Li, T. & Katz R.S. (2010). EQ-5D and SF-36 Quality of Life Measures in Systemic Lupus Erythematosus: Comparison with Rheumatoid Arthritis, Non-inflammatory Rheumatic Disorders, Fibromyalgia. *The Journal of Rheumatology*, 37, 296-304.
- Wolfe, F., Smythe, H.A., Yunus, M.B., Benett, R.M., Bombardier, C. et al. (1990). The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the Multicentre Criteria Committee. *Arthritis, Rheumatology*, 33, 160-172.
- Woolf, C.J. (2010). Central sensitization: Implications for the diagnosis and treatment of pain. *Pain*, 152, 2-15.
- Yunus, MB. (2002). Gender differences in fibromyalgia and other related syndromes. *The Journal of Gender-Specific medicine*, 5, 42-47.

Journal Article 2

The relationship between a two factor model of self-compassion, self-esteem and pain in a chronic pain population

Authors

Caitríona Doughty, Dr. Laura Keith, Dr. Paul G. Morris, & Dr. Nuno Ferreira

Affiliations

Department of Clinical Psychology NHS Fife, Clinical and Health Psychology, University of Edinburgh Scotland.

Corresponding Author

Caitríona Doughty, Clinical Psychology Department, Lynebank Hospital, Halbeath Road, Dunfermline, Fife, KY11 4UW

Written in accordance with the instructions for '*The Clinical Journal of Pain*' (see Appendix A for author guidelines)

Abstract

Objectives: This study aims to explore the predictive role of self-compassion (self-warmth and self-coldness) and self-esteem in key outcomes of quality of life and disability in a chronic pain population.

Methods: A quantitative cross-sectional design was used. Sixty participants, recruited through NHS chronic pain services and pain self-management groups, completed measures of pain severity, self-compassion, self-esteem, distress, quality of life and disability. Correlation and hierarchical regression were used to explore relationships between these variables.

Results: Quality of life was lower than previous research, with females reporting higher levels of quality of life than males. After controlling for demographic and clinical variables, self-warmth was a significant predictor quality of life, however it was not predictive of disability. Neither self-coldness nor self-esteem were significant predictors of quality of life or disability.

Discussion: Self–warmth is associated with better quality of life in people with chronic pain. Psychological interventions targeting the development of self-warmth attributes and skills may have beneficial effect in the psychological adjustment of people suffering with chronic pain. Further research is needed to replicate these findings with a larger and more representative sample.

Keywords: self-compassion, self-esteem, self-coldness, self-warmth, quality of life, chronic pain,

2.1 Introduction

Chronic pain has been defined by the Scottish Intercollegiate Guidelines Network as the experience of pain for at least three months that does not resolve with treatment¹. With a worldwide estimated prevalence of 19-34%,^{2,3} chronic pain is estimated to be one of the conditions with greatest socio-economic costs.⁴ As the burden of chronic pain increases, predictors and outcomes of chronic pain have received growing attention⁵. Psychological factors are widely accepted as influencing the relationship between chronic pain and outcomes such as disability and distress.⁶ Psychological approaches have been applied in chronic pain management and seem beneficial.⁷ However, psychological treatments for chronic pain have shown small effect sizes and further development is required⁸. It has been suggested that a focus on understanding the treatment mechanisms or processes may help to increase the efficacy of these treatments.⁹ One set of processes that have started to receive growing attention in the area of chronic pain are those related to self or personal identity such as self-compassion and self-esteem.¹⁰

2.1.1 Evidence for Self-Compassion as a Predictor

It has been suggested that positive affect may provide a buffering function to health problems associated with negative emotions in patients with chronic illness¹¹. The role of self-compassion was explored in a chronic illness group and a cancer group in relation to psychopathology and quality of life¹². Pinto-Gouveia et al. (2013) used a two-factor model of the Self-Compassion Scale referred to as self-coldness and self-warmth and it was found that self-coldness was the best predictor followed by self-warmth for quality of life in the chronic illness group. However Pinto-Gouveia et al. (2013) also found that self-warmth was the best predictor of quality of life for individuals with cancer. Pinto-Gouveia et al.

(2013) study reported limited demographic and clinical information and controlled only for education level, therefore it was difficult to infer what factors may have influenced the differences between the groups e.g. gender, age, severity of pain. Several studies have revealed that pain severity and distress are factors that contribute to poor quality of life and disability.^{13,14} Participants in Pinto-Gouveia et al. (2013) study were predominantly female and middle aged. A recent meta-analysis by Yarnell et al., (2015) suggested that while gender differences on levels of self-compassion were significant, only a small effect size was observed.¹⁵ Neff and Pommier (2013) proposed that women are often assigned the role of caregiver in society, however they are less able than men to relate to themselves with the same level of compassion that they would show to others.¹⁶ It was also found that females tend to be more critical of themselves and demonstrate more negative self-talk than males. Age has been found to reduce the effect of gender differences on levels of self-compassion.¹⁵ Ardelt et al (2010) argued that the understanding of common humanity is a wisdom that is more likely to increase with age regardless of gender because people have more life experience to reflect upon.¹⁷

2.1.2 How Self-Compassion is conceptualised and measured

Self-compassion has been defined as comprising three main components: 1) self-kindness in the absence of self-judgement or self-criticism; 2) common humanity whereby an individual perceives their struggle as part of a shared human experience rather than an isolated experience; 3) mindfulness without over-identification in relation to negative thoughts or feelings. It is argued that when these components interact, an individual has the tendency to be caring, warm and understanding towards oneself regardless of one's successes or failures ¹⁸.

Self-compassion has been suggested to improve awareness and emotion regulation associated with psychopathology¹⁹⁻²¹. MacBeth et al. (2012) conducted a meta-analysis on the relevance of self-compassion for the emotional symptoms of anxiety, depression and stress.²¹ Findings showed that self-compassion is related to psychopathology, identifying a large effect size. Associations were reported to be significant regardless of clinical status, age or gender. This meta-analysis used the total score from Self-Compassion Scale (SCS)²² and the Self-Compassion Scale-Short Form (SCS-SF)²³ rather than specific subscale scores. A recent meta-analysis by Muris et al (In Press) focused on the subscales of the SCS and SCS-SF in relation to psychopathology.²⁴ It was found that the associations between the negative subscales of the SCS (self-judgement, over-identification, isolation) and psychopathology were significantly stronger than the positive subscales of the SCS (self-kindness, common humanity and mindfulness). This suggests that the use of a total compassion score of the SCS or SCS-SF, which includes the reversely scored negative subscales, will potentially result in an inflated association with symptoms of psychopathology. This would result in an overestimation of the presumed protective role of self-compassion in mental health problems. Including both positively and negatively worded items in questionnaires were introduced with the aim of reducing response bias, however studies have shown that it can potentially undermine the quality of the assessment.²⁵ This may be the case for the SCS as a recent factor analysis did not support its construct validity when using a total score.^{26, 27} For this reason, some researchers into self-compassion have started separating the positive (self-kindness, common humanity, mindfulness) and negative (self-judgement, isolation, over-identification) aspects of self-compassion into two dimensions known as self-warmth and self-coldness (Gilbert, McEwan, Matos, & Rivis, 2011).²⁸

It has been suggested that to clarify the protective nature of self-compassion, it would be appropriate to discard self-coldness (self-judgement, isolation and over-identification) from the questionnaire and to focus solely on self-warmth (self-kindness, common humanity and mindfulness).¹⁵ In addition, compassion focused interventions have reported specific improvements in positive affect, but they do not seem to effectively reduce negative affect more than any other intervention ^{28, 29}. This conclusion may be explained as a lack of large-scale, high-quality trials rather than the existence of negative evidence.³⁰

Although there is evidence to suggest that self-compassion is associated with increased motivation in relation to positive health behaviours,³¹ this element is not represented in the items of the SCS or SCS-SF. As a result, the three aspects of self-warmth (self-kindness, common humanity and mindfulness) are not necessarily distinctive of compassion but seem to tap a number of already known protective factors. For example, there is a vast amount of research evidencing that mindfulness on its own promotes well-being in chronic pain populations. ³²

Constructs such as belongingness ³³ and self-soothing ³⁴ closely resemble Neff's ¹⁸ dimensions of common humanity and self-kindness. It would be useful to explore which components of self-compassion are important within the context of psychopathology. ²⁴ This is also true for the relative importance of self-compassion as compared with other constructs that have been advocated in the positive psychology literature such as self-esteem.²⁴ Although global self-esteem is considered to be less stable and more reactive than self-compassion,³⁵ both concepts are viewed as highly relevant for a person's general feeling of self-worth and as such it is not surprising that their link has often been studied

in previous research.^{81,35-37} Thus, in order to study the unique relationship between self-compassion and mental health problems, it seems important to control for global self-esteem.

2.1.3 Overlap with other concepts

High self-esteem has been found to be a resource for coping with illness,³⁸ buffering against depression, stress, and negative emotions.^{39,40} In particular, self-esteem has been linked to psychological adjustment in various chronic diseases such as cystic fibrosis, multiple sclerosis and HIV.⁴¹⁻⁴³ Conversely, illness may impact negatively on self-esteem.⁴⁴ Self-esteem may be threatened by experiencing illness-related consequences, or by decreased coping with everyday life challenges.^{45,46} In turn, low self-esteem may lead to unhealthy behaviours in an attempt to self-regulate negative emotions.⁴⁷ A previous longitudinal study suggests that low self-esteem causes more negative affect for chronic disease patients than healthy populations.³⁹ As self-esteem can be an important coping resource, one of the concerns raised by the combination of low self-esteem and chronic illness is an increased difficulty in preventing the negative affect from developing into a more severe case of depression. Research demonstrated that patients with chronic pain who had lower self-esteem reported more negative affect, less positive affect and greater symptom severity than individuals with higher self-esteem.³⁸

Neff⁴⁸ distinguished self-compassion from the related construct of self-esteem by proposing that self-esteem is based on evaluating oneself in comparison with others whereas self-compassion is based on relating to oneself without self-evaluation or comparison. Despite the benefits of self-esteem, studies show that what people do to get and maintain a sense of high self-esteem is linked to negative outcomes such as narcissism

and increased aggression.^{49,50} Reasons for this may be that self-esteem is often dependent on favourable comparison with others and successful attainment which results in fluctuation depending on one's successes or failures.⁴⁸

A number of studies have been conducted to investigate the cognitive and emotional processes by which self-compassionate individuals manage negative life events.⁵¹ It was evidenced that self-compassion accounted for unique variance in the outcome variables when controlling for self-esteem. It was also reported that because self-compassion is not evaluation based, self-compassion appears to be a more stable and unconditional form of self-regard than self-esteem.³⁵ It must be noted that the situations the participants were dealing with in these studies⁵¹ were unchallenging such as receiving a poor grade on a test or neutral feedback in comparison to many of the hardships people encounter in daily life. Therefore, their research shows that being self-compassionate helps people deal with real and imagined negative events, but the question remains of the degree to which treating oneself kindly buffers people against the impact of severe challenges in everyday life such as chronic pain.

In a study examining resilience levels in adults with congenital spinal bifida, it was found that chronic pain negatively impacted on self-esteem and self-compassion. This is in line with previous research in that disability severity can result in frustration and distress.⁵² The findings that self-esteem and self-compassion correlate with resilience are consistent with the idea that having a source of positive self-regard and an ability to be kind to oneself is likely to promote positive adaptation when faced with adversity. A strong correlation was found between self-esteem and self-compassion ($r=.688$) which is in line

with previous findings ^{18,51,53}, raising the question of whether these are psychological distinct constructs.

2.1.4 Summary

Research in chronic pain has tended to focus on specific diagnoses, such as neuropathic pain ⁵⁴ and fibromyalgia,⁵⁵ or on specific subgroups such as older people ⁵⁶ and trauma survivors.⁵⁷ This focus on specific diagnoses/subgroups is limiting because many people with chronic pain do not have a clear diagnosis or precipitant for their pain.⁵⁸ For this reason, the present study did not focus on one particular pain condition.

Given the growing interest in the association between emotion regulation problems and unhelpful forms of self-relating with chronic pain, understanding the role of self-compassion and self-esteem may be potentially important variables in chronic pain in order to develop more effective treatment interventions. Muris et al. (In press) conducted a meta-analysis which found a significant link between self-warmth, self-coldness and mental health indicators.²⁴ Self-compassion also predicts unique variance in mental health after controlling for self-esteem. It was suggested that it would be useful to explore whether a two factor model of the SCS (self-warmth and self-coldness) is a better fit as this model seems to have stronger psychometric validity in both clinical and non-clinical samples.⁵⁹ A recent review of the literature indicated that the vast majority of studies (83.5%) relied only on the SCS or SCS-SF total score as an index of self-compassion.²⁴ Studies including self-esteem, self-warmth, self-coldness as predictors are limited in a clinical population and non-existent in a sample of participants with chronic pain. Previous literature has indicated that demographic variables such as age, gender and pain duration contribute to levels of self-compassion and pain outcomes including quality of life and

disability.^{15,60} For this reason, the current study will control for relevant demographic variables when investigating self-warmth, self-coldness and self-esteem as predictors of disability and quality of life.

2.1.5 Aims

The main aim of the study was to determine whether and to what extent self-warmth, self-coldness and self-esteem would be unique predictors for pain outcomes while controlling for relevant demographic and clinical variables. We predicted that self-warmth and self-esteem would be (positively) associated with quality of life and (negatively) associated with disability, severity of pain and distress. We predicted that self-coldness will be (negatively) associated with quality of life and (positively) associated with disability, severity of pain and distress. It was hypothesised that self-warmth and self-coldness would be stronger predictors than self-esteem for pain outcomes, disability, and quality of life after controlling for known predictors such as demographic (e.g. gender and age) and clinical variables (e.g. distress and severity of pain)

2.2 Method

2.2.1 Design

This study used a cross-sectional, questionnaire-based design to explore relationships between self-warmth, self-coldness, self-esteem, and pain outcomes (distress, disability and quality of life). Ethical approval was granted by the NHS South East Scotland Research Ethics Committee (See appendix K).

2.2.2 Participants

During the period of February 2015 to February 2016, four hundred participants were invited to participate from the NHS Chronic Pain Service in Fife and Lothian, the NHS Fife Rheumatoid Disease Unit and the Fife Pain Association Scotland Self-Management Groups. Sixty individuals returned completed questionnaires (return rate= 15%). Forty percent of the responses were returned from the Chronic Pain Service in Fife and Lothian, eighteen percent from the NHS Fife Rheumatoid Disease Unit and forty-two percent from the Fife Pain Association Scotland Management Groups. Non-identifiable demographic information was collected in order to establish the characteristics of the sample. A total of 39 individuals (65%) were female and 21 individuals (35%) were male. Their mean age was 53.08 years (SD = 14.49) and their mean duration of pain was 11.2 years (SD= 10.7). Mean self-reported pain intensity and pain interference measured by the Brief Pain Inventory- Short Form (BPI-SF),⁶¹ was 6.46 out of 10 (SD= 1.59) and 6.92 out of 10 (SD= 2). These pain scores for the BPI-SF subscales were similar to reported scores on the BPI-SF in other studies carried out in a chronic pain.⁶² Participants had a variety of pain sites, with joint pain being the most common (34%), followed by back pain (28%), all over body pain (17%), and right sided pain (7%). The remaining pain locations were neck and leg. They reported having had a variety of treatments, with prescribed medication, physiotherapy and reflexology being the most commonly reported treatments.

2.2.3 Measures

Permissions for the use of copyrighted questionnaires were received prior to administration (Appendix I)

Demographics Questionnaire: Participants were asked to provide non-identifiable demographic data including age, gender, pain duration, diagnosis, and who provided their diagnosis.

Self-Compassion Scale: (SCS)²² contains 26-items assessing how people typically act towards themselves in adverse circumstances. There are three factors of positive self-compassion: self-kindness; common humanity; and mindfulness, and three factors that focus on lack of self-compassion: self-judgement; isolation; and over-identification. This study obtained two separate scores for this scale: self-warmth (sum of three positive factors) and self-coldness (sum of the three negative factors). Responses to each item are measured on a five-point scale from “Almost never” to “Almost always”. The scale has good predictive, convergent and discriminant validity and has been shown to have good internal consistency when used with a pain population (0.91-0.95).^{60,63} Cronbach’s alpha for the Self-warmth subscale in this sample was .92. Cronbach’s alpha for the Self-coldness subscale in this sample was .91.

The Rosenberg Self-Esteem Scale: (RSES)⁶⁴ includes 10 statements related to feelings of self-worth or self-acceptance, and measures of global self-esteem. The response set for this scale was 1 (strongly agree) to 4 (strongly disagree) with total scale scores ranging from 10 (low self-esteem) to 40 (high self-esteem). The scale has good predictive, convergent and discriminant validity and ⁶⁵ and has been shown to have good internal consistency when used in a clinical population for individual with HIV and Cystic Fibrosis with a Cronbach’s alpha ranging from 0.85 to 0.94. ^{43,66} Cronbach’s alpha for the RSES in this sample was .88.

Brief Pain Inventory- Short Form: (BPI-SF)⁶¹ comprised 15 items and assesses for the presence of pain, pain severity and functional interference from pain (general activity, mood, walking, relationships with others, sleep, normal work and enjoyment of life) in the last 24 hours. For these subscales a higher mean score indicates higher pain severity and pain interference. It also assesses the types of pain medications being used, the percentage of pain relief obtained from medications and considers the distribution of pain via a body map. The BPI-SF has been used extensively in clinical trials for pain and has been recommended as a reliable and valid measure.⁶⁷ Internal consistency for the pain severity score and for the interference scale has been reported as ranging from 0.85 and 0.88 respectively.⁶⁸ Construct validity of the BPI-SF has been supported for the generic assessment of pain as well as specifically for low back pain and rheumatoid arthritis.⁶⁹ Cronbach's alpha for the BPI-SF in this sample was .91.

The Hospital Anxiety and Depression Scale: (HADS)⁷⁰ is widely used for measuring anxiety and depression in clinical population with physical illness. It contains 14 items and requires the respondent to rate the degree to which they have experienced emotions relating to distress in the past week. This scale provides an overall measure of emotional distress as well as depression and anxiety subscale scores. All items are rated on a four-point response scale (0-3). Cut off scores for both anxiety and depression are normal (0-7), mild (8-10), moderate (11-15) and severe (16-21).⁷⁰ A person with a total score of 15 or more (15+) on the HADS is considered to experience clinically significant emotional distress.⁷¹ Across a wide range of samples the reliability and validity of the HADS has been well established with a Cronbach's alpha for each scale ranging between .67 and .93.^{72,73} Cronbach's alpha for the HADS in this sample was .91.

Pain Disability Questionnaire: (PDQ)⁷⁴ incorporates a disability-related psychosocial component in addition to a physical functioning component related to pain. The psychosocial component consists of nine items and has a score range of 0-90, which measures how much pain affects mood, physical role, leisure activities. The functional status component contains six items and scores range from 0-60 and measures how much pain impacts on employment status, self-care, mobility and income. The PDQ yields a total functioning disability score ranging from 0 (optimal function) to 150 (total disability), using 10 point Likert scale.⁷⁵ The PDQ has good reliability with a Cronbach's alpha of .96. The construct related validity of the PDQ was found to be excellent as it correlated well to both the Million Visual Analog Scale (0.65-0.81) and Oswestry Pain Disability Questionnaire (0.55-0.80).⁷⁴ Cronbach's alpha for the PDQ in this sample was .92.

Quality of Life Scale: (QOLS)⁷⁶ is a 16 item questionnaire that measures physical and mental well-being, relationships with other people, social, community and civic activities, personal development and fulfilment, recreation and independence. Each item was scored 1-7 and the total score ranges from 16-112. A higher total score shows a higher QOL. This scale has shown good validity and test-retest reliability.⁷ Good internal consistency ($\alpha=0.88-0.93$) was found in a number of studies with fibromyalgia participants and a coronary obstructive pulmonary disease population.^{78,79} Cronbach's alpha for the QOLS in this sample was .93.

2.2.4 Procedure

NHS professionals in the Pain Management and Rheumatoid services, and facilitators in the Pain Association Self-Management Groups provided people attending these services with packs containing information about the study (Appendix L) and the six research

questionnaires (Appendix N) along with a general questionnaire collecting non-identifiable demographic information (Appendix M). Individuals who were interested in taking part in the study were given the packs to take home which they could then post back to the researcher using an included pre-paid envelope. Consent to participate was provided by the return of the (anonymous) research questionnaires. The questionnaires were marked with a unique identifier code for each recruitment site.

2.2.5 Statistical Analysis

All statistical analyses were conducted using IBM SPSS Statistics Version 22. The computational and modelling tool PROCESS⁸⁰ for SPSS was used for mediation analyses.

2.2.6 Power and Sample Size

Power analyses were conducted to guide recruitment. As the relationships between self-compassion, self-esteem and pain outcomes (disability and quality of life) had not previously been investigated in this population, a medium effect size was assumed (conservatively), based on previous studies between self-compassion and general psychopathology in adults ($r = -0.54$; 95% CI $[-0.57, 0.51]$)²¹ and chronic pain population (r 's ranging from -0.50 to -0.52)^{73,60}. It was determined that in order to have .8 power to detect a medium effect size at an alpha level of .05 with six independent variables (gender, severity of pain, distress, self-warmth, self-coldness and self-esteem) and two dependent variables (disability, quality of life) a sample of 103 was required (G*Power Version 3.0.10).

2.2.7 Missing Data

Data was checked for missing items. Little's Missing Completely at Random (MCAR) Test was found to be non-significant ($\chi^2(72) = 65.74, p < .69$). As data was found to be MCAR, missing data was dealt with by using individual mean substitution.⁸² This meant that for measures where up to one item per subscale was missing but less than 10% of total scale, and where there was homogeneity within subscales, individual mean substitution was adopted i.e. the inputted value was calculated from the mean of the available items of that subscale/total scale for the given respondent. While internal consistency can be positively skewed by using this method, it does allow for data to be included where limited items are missing. A total of 10 cases had missing items replaced using this method.

2.2.8 Tests of Reliability

Cronbach's alpha tests of internal consistency were calculated for key variables and subscales. Values above .7 are generally considered to suggest adequate levels of reliability.⁸³ Within this study reliability for all subscales and scales ranged from 0.76 to 0.93.

2.2.9 Data Analysis

Data was initially screened to ensure that assumptions of further analyses were met. Histograms and boxplots were examined to ensure no outliers were present. Assumptions of linearity and homoscedasticity were found to be met through examination of scatterplots. Pearson correlations were calculated between all predictor variables to test for multicollinearity. No extremely high correlations i.e. 0.9, were identified, suggesting that all items were suitable for inclusion in further analyses.⁸³ The Shapiro-Wilk test at $p > 0.05$ was used to test for normality based on recommendations by ⁸³ for small sample sizes.

Independent t-tests were used to explore gender differences. Correlational analysis was used to examine how the demographic (age) and clinical variables (severity of pain, pain duration and distress) related to self-warmth, self-coldness, self-esteem and pain outcomes (quality of life and disability) to establish what variables to control for in the hierarchical regression analyses. Correlational analyses were also run to determine the relationship between self-warmth, self-coldness, self-esteem and the outcome variables (quality of life and disability). Following this hierarchical regression analyses using forced entry of three blocks were planned with self-warmth, self-coldness and self-esteem as the predictor variables and disability and quality of life as the dependent variables while controlling for gender, pain severity and distress.

2.3 Results

2.3.1 Descriptive Data

Table 2.1 presents the difference between males and females in relation to mean and standard deviation (SD) for each measure completed in the study. For the total samples self-esteem scores produced a mean of 26 (SD 7.2). Average scores on the SCS subscales were 35.34 (SD= 11.68) for self-warmth and 43.78 (SD=11.65) for self-coldness. Mean scores on the HADS subscales were 11.56 (4.87) for anxiety and 9.88 (4.67) for depression and when classified according to clinical thresholds produced, 70% (N=42) were in the clinical range of symptoms for anxiety and 80% (N=48) for depression. Mean scores of 63.67 (SD=18.58) and 93.12 (SD=26.61) were obtained for quality of life and disability respectively.

2.3.2 Control Variables

Gender has been shown to be a significant influence on many aspects of life for adults with chronic pain.⁸⁴ Preliminary two-tailed t-tests were conducted to compare scores of males (N=21) and females (N=39) on each measure (see Table 2). Females reported higher quality of life with the Cohen's d indicating a medium effect size (0.7). No significant gender differences were found for levels of distress, pain severity, self-esteem, self-warmth and self-coldness. Cohen's d indicated small effect sizes for these non-significant mean differences. Significant gender differences found in relation to quality of life indicated that gender should be controlled for in the hierarchical regression where quality of life was the dependent variable.

Table 2.1 Mean differences between genders on outcome measure scores

Outcome Measures	Male Mean (SD) N=21	Female Mean (SD) N=39	Mean Difference	t	P-value	95% Confidence Interval Lower Upper	
BPI-SF Pain Severity Subscale	6.38 (1.6)	6.50 (1.6)	.119	.28	.79	-.76	.99
HADS TOTAL SCORE	23.95 (7.3)	20.10 (9.0)	3.85	-1.79	.08	-8.18	.48
Rosenberg Self-Esteem Total Score	25.38 (6.1)	26.48 (7.8)	1.20	.60	.55	-2.56	4.76
Self-Warmth	34.52 (12.8)	35.77 (11.2)	1.25	.38	.71	-5.48	7.98
Self-Coldness	43.79 (12.0)	43.77 (11.6)	-.02	-.00	.99	-6.52	6.49
Quality of Life Total Score	55.61 (17.3)	68.00 (18.0)	12.39	2.61	.01*	2.80	21.98
Pain Disability Questionnaire Total Score	101.92 (22.2)	88.34 (27.8)	-13.54	-1.92	.06	-27.63	.56

*Gender difference significant at the level of $p < .05$

95% bias corrected and accelerated confidence intervals based on 1000 bootstrap samples

*Levene test significant

2.3.3 Correlation Analysis

Table 2.2 presents the results of the Pearson's correlations among the measures of self-warmth, self-coldness, self-esteem, severity of pain, quality of life, distress and disability. All variables were significantly correlated in the predicted directions. Consideration was given to the inclusion of age and duration of pain as control variables within the regression analysis. However, age did not significantly correlate with either quality of life [CI-.14,.39] or disability [CI-.5,.08] and pain duration did not significantly correlate with either quality of life [CI-.19, .33] or disability [CI-.31,.29].

2.3.4 Regression Analysis

A four stage hierarchical regression was conducted with quality of life as the dependent variable, to explore the independent effect of self-esteem, self-warmth and self-coldness on quality of life while controlling for gender, severity of pain and distress. Data met the assumption of independent errors (Durbin-Watson= 1.9) and multi-collinearity was not deemed to be a concern as tolerance scores ranged from .29 to .8 and variance inflation factor (VIF) scores from 1.26 to 3.48 (O'Brien, 2007). When gender was entered in step 1, it explained 8.7% of the variance in quality of life. Entering severity of pain subscale at step 2 explained an additional 13% of the variance in quality of life. HADS total score was entered in at step 3 which explained 42% of the variance in quality of life. Finally, RSES total score and the SCS subscales (self-warmth and self-coldness) were entered in at step 4, explaining 8% of the variance in quality of life. The model as a whole explained 70.6% of the variance in quality of life. Four variables made a statistically significant contribution to this model. In order of highest beta values: HADS total, self-warmth subscale, and BPI-SF pain severity. The negative coefficients indicate that higher levels of distress, and pain

severity were associated with lower rates of quality of life. The positive co-efficient indicates that higher levels of self-warmth were associated with higher rates of quality of life. Coefficients for all variables, including those with a non-significant contribution to the model, are presented in Table 2.3.

A second hierarchical regression analysis was performed to examine if self-warmth, self-coldness and self-esteem made a unique contribution to disability outcomes after pain severity, and distress were accounted for in the analysis. BPI-SF pain severity subscale was entered in step 1 and explained 18% of the variance in disability. Entering HADS total score at step 2 explained an additional 17% of the variance in disability. RSES total score and the SCS subscales self-warmth and self-coldness were entered in at step 3. RSES total score and self-compassion subscales were not significant predictors of scores on the PDQ.

Table 2.2 Pearson's correlation coefficients for correlations of key variables, age and pain duration

	1.	2.	3.	4.	5.	6.	7.	8.
1. Rosenberg Self-Esteem Score								
2. Self-Warmth	.482**							
3. Self-Coldness	-.600**	-.435**						
4. HADS Total	-.661**	-.485**	.617**					
5. Quality of Life Scale	.584**	.603**	-.496**	-.786**				
6. Pain Disability Questionnaire	-.519**	-.349**	.398**	.538**	-.649**			
7. BPI-SF Severity Subscale	-.281*	-.015	.162	.336**	-.344**	.436**		
8. Age	.235*	.227*	-.150	-.261*	.124	-.214	-.056	
9. Pain Duration	.012	.039	-.346**	-.231*	.083	-.038	-.118	.294*

Based on 2000 bootstrap samples

** Correlation is significant at the 0.01 level (one-tailed), * Correlation is significant at 0.05 level (one-tailed)

Table 2.3 Summary of Hierarchical Regression to predict quality of life

Variable	β	95% CI		B	t	p-	R	R ²	ΔR^2
		Lower	Upper	standardised		value			
Step 1							.32	.10	.09
Gender	-12.39	-22.01	-2.77	-.32	-2.58	.01			
Step 2							.48	.23	.20
Gender	-12.89	-21.89	-3.89	-.33	-2.87	.01			
BPI-SF Pain Severity	-4.17	-6.91	-1.44	-.36	-3.06	.00			
Step 3							.81	.65	.63
Gender	-6.64	-12.93	-0.35	-.17	-2.11	.04			
BPI-SF Pain Severity	-1.30	-3.28	0.68	-.11	-1.31	.19			
HADS Total	-1.53	-1.91	-1.16	-.71	-8.25	<.00			
Step 4							.86	.74	.71
Gender	-7.63	-13.37	-1.89	-.20	-2.67	.01			
BPI-SF Pain Severity	-1.96	-3.79	-0.13	-.17	-2.15	.04			
HADS Total	-1.10	-1.59	-0.62	-.51	-4.59	<.00			
Self-Warmth	.528	.25	.80	.33	3.88	<.00			
Self-Coldness	.013	-.30	.32	.01	.08	.93			
RSES	.075	-.46	.61	.03	.28	.78			

95% bias corrected and accelerated confidence intervals based on 1000 bootstrap sample

Table 2.4 Summary of Hierarchical Regression to predict pain disability

Variable		β	95% CI		B standardised	t	p-value	R	R ²	ΔR^2
			Lower	Upper						
Step 1								.44	.19	.18
BPI-SF	Pain	7.32	3.34	11.29	.44	3.69	<.00			
	Severity									
Step 2								.60	.36	.34
BPI-SF	Pain	4.82	1.05	8.60	.29	2.56	.01			
	Severity									
HADS Total		1.36	.668	2.06	.44	3.93	<.00			
Step 3								.64	.41	.36
BPI-SF	Pain	4.96	1.12	8.79	.30	2.59	.01			
	Severity									
HADS Total		0.67	-0.30	1.65	.22	1.38	.17			
Self-Warmth		-0.27	-0.85	.30	-.12	-.95	.35			
Self-Coldness		0.08	-0.57	.73	.04	.25	.80			
RSES		-0.79	-1.9	.34	-.21	-1.4	.17			

95% bias corrected and accelerated confidence intervals based on 1000 bootstrap samples

2.4 Discussion

This cross sectional study aimed to explore the role of self-warmth, self-coldness and self-esteem as predictors of quality of life and disability in a chronic pain sample. The results of the study will now be considered in relation to the initially outlined aims and hypotheses. Results showed that participants' levels of self-esteem,⁸⁵ pain severity,⁶² disability,⁷⁴ and distress⁸⁶ were in line with previous studies using a chronic pain sample. Quality of life was slightly lower in this study than in previous studies^{77,87} and it was not possible to compare the self-warmth and self-coldness subscales as they have not been specifically assessed in a chronic pain sample. Mental health symptoms were at clinical level for 77% of the sample, which is consistent with other studies indicating elevated mental health difficulties in a chronic pain population.⁸⁸ Overall these scores suggest that the study sample was representative of a chronic pain population. One potential reason for the lower overall mean levels of quality of life in this study was the significant gender difference between male and females on this measure. Gender differences on levels of quality of life in a chronic pain sample remain inconclusive as a number of studies have found males to have higher quality of life⁸⁹ while others⁸⁷ did not confirm any significant differences. Rustoen⁹⁰ reported similar results to this study where they found that females reported higher quality of life compared to males. One possible explanation for higher quality of life scores in women in this sample may be that men and women cope differently with chronic pain⁹¹. While men tend to use behavioural distraction and problem-focused tactics to manage pain, women tend to use a range of coping techniques including social support, positive self-statements, emotion-focused techniques, cognitive reinterpretation, and attentional focus.⁹²

Correlation analyses revealed that higher self-esteem, higher self-warmth and lower self-coldness were related to lower levels of distress, lower disability and better quality of life.^{60,38} It was found that age was significantly related to higher self-esteem, self-warmth, and lower distress. This is in line with other studies which suggest that self-compassion on well-being has been shown to be more beneficial to middle-aged adult than those in early adulthood.⁹³ It may be that an understanding of common humanity increases with age.¹⁷

The significant relationship between higher pain severity, higher distress, higher disability and lower quality of life are consistent with the results of previous research.^{6,12} The significant relationships found within this study were in contrast to previous findings in a chronic illness group whereby self-compassion and self-esteem did not correlate significantly with lower disability.⁵² Our findings indicate that having a salient source of positive self-regard and an ability to be kind to oneself is likely to promote positive adaptation when faced with adversity. Previous literature has reported the strong correlation between self-esteem and self-compassion, suggesting that the two constructs are not psychologically distinct. This study found that while self-coldness strongly correlated (-.6) with self-esteem, self-warmth only demonstrated a medium correlation (.42) with self-esteem. Muris et al. (in press)⁹⁴ reported a similar correlation between self-warmth and self-esteem (.44). They noted that common humanity which is a component of this subscale did not correlate significantly with global self-esteem, possibly because self-esteem is more concerned with comparing oneself to others rather than feeling connected to others. Nevertheless, the correlation between self-warmth, self-coldness and self-esteem were in keeping with what has been reported in the literature

³⁵ and the magnitude of the inter-correlations demonstrate that self-warmth, self-coldness and self-esteem appear to be distinct but related to aspects of personal identity. ⁹⁵

Hierarchical regression revealed that self-warmth was a significant predictor of quality of life after controlling for gender, pain severity and distress. Self-coldness was not found to be a significant predictor for quality of life. These findings were in line with previous research carried out among patients with chronic illness and cancer.¹² These studies found that self-warmth was the only predictor for quality of life among cancer patients. However, Pinto-Gouveia et al., 2013 also reported that self-coldness explained the majority of the variance for quality of life in patients with chronic illness.¹² It would be expected that the results of this study should be more in line with the chronic illness patient group rather than the cancer patient group in Pinto-Gouveia et al. (2013) study.¹² However, their study did not contain a pain severity measure and therefore it may be that the cancer patients experienced more pain severity ⁹⁶ than the chronic illness group, which might account for the similarities between the chronic pain sample in the present study and the cancer group in Pinto-Gouveau et al. (2013) study.¹² These findings seem to support the theory that the way self-compassion enhances positive wellbeing is via the increased self-kindness, common humanity, and mindfulness associated with a compassionate mind state. It must be noted that distress was the more important predictor of quality of life than the other predictors in the model. Distress was added to the forced entry regression before self-coldness and it is likely that this reduced the predictive power of self-coldness in this model. This study and previous studies have reported significantly high correlations between self-coldness and distress compared with self-warmth and distress. This finding offers further support to Muris et al (2016) argument

that although the protective influence of self-warmth exists in relation to psychopathology, the effect is likely to be boosted when self-coldness is also included.²⁴

The hypothesis that self-esteem would be a significant predictor for quality of life was not supported in this study. The finding may be a result of the measure used for quality of life in that the QOLS does not contain subscales separating out physical and psychological components of quality of life like the SF-36.⁹⁷ In a study looking at multiple sclerosis, it was found that self-esteem only explained the variance on the Mental Component Scale (MCS) rather than the Physical Component Scale (PCS) of the SF-36.⁴² Self-esteem thus seems to be important in MCS, which is in line with previous findings in other chronic conditions.^{43,85,98} Many studies used the SF-36 in medical setting for chronic illness or chronic pain and it may be that self-esteem is more likely to predict quality of life because self-esteem is a psychological variable and MCS may be considered predominantly of psychological or emotional nature. While self-esteem was not a predictor for quality of life, our findings support previous findings which reported that self-compassion would account for the unique variance in outcome variables when controlling for self-esteem. This study adds to the evidence base in that chronic pain is a more adverse life experience than the examples (e.g. poor grade on a test) that were used in other research when examining the differences between self-esteem and self-compassion in relation to psychological well-being.⁵¹

The findings that neither self-warmth, self-coldness nor self-esteem were significant predictors of physical disability were contrary to our hypotheses that self-warmth or self-coldness would be a stronger predictor than self-esteem for disability. In contrast other research demonstrated that self-warmth predicted better adjustment to physical disability.⁶⁰

It must be noted that Wren et al.⁶⁰ recruited participants who had a >30 BMI, therefore it may be that self-warmth has the potential to act as a buffer against the mental and global health detriments of self-stigma which relate to disability in overweight and obesity⁹⁹ rather than chronic pain. Our study replicated findings from Hoge et al.¹⁰⁰ in that they found that self-compassion correlated with disability in a group of people suffering from generalised anxiety disorder, however, self-compassion was not a significant predictor of disability. Instead they found that the non-judgment of inner experience component of mindfulness contributed to lower disability. This has also been reported in the chronic pain literature³² and it may be useful to further explore the subcomponents of self-warmth (common humanity, self-kindness and mindfulness) to explore which components correlate and predict a relationship with lower disability in chronic pain.

2.4.1 Clinical Implications

The current findings have important implications for clinicians working with this population. Clinicians should assess psychological constructs such as self-warmth and target interventions to promote this type of protective processes in an attempt to improve overall quality of life in people with chronic pain. There is rising evidence that self-warmth can be increased with practice¹⁰¹ with many interventions having been proposed for people with chronic pain; loving kindness meditation;¹⁰² compassion meditation;¹⁰³ positive psychology techniques;¹⁰⁴ and mindfulness.³² There is also research to suggest that third wave interventions such as Dialectical Behaviour Therapy (DBT)¹⁰⁵ and Acceptance and Commitment Therapy (ACT)¹⁰⁶ may be useful due to their emphasis on mindfulness and concepts similar to over-identification i.e. cognitive defusion.¹⁰⁷ In keeping with recent government strategies which

state that “good mental health is not potentiated solely by the absence of mental ill health but the presence of positive mental health factors”,¹⁰⁸ it is suggested that increased knowledge on the role of self-compassion in adult mental health, particularly chronic pain settings, may provide and promote an alternative way to conceptualise chronic pain difficulties.

2.4.2 Strengths and Limitations

To the author’s knowledge, this study is the first to explore the role of self-warmth, self-coldness and self-esteem as predictors of quality of life and disability in a chronic pain sample. The SCS is currently the most widely used measure of self-compassion, however concerns have been raised in relation to its underlying structure. Costa et al. (2015) did not replicate evidence for an overarching construct of self-compassion, suggesting that a two factor structure model is more appropriate.⁵⁹ The present study used a two factor model to further investigate its utility in a clinical population. In addition, the study used standardised validated psychometrics with psychometric properties demonstrated both in previous research and replicated with the current sample. The sample was recruited from several sites in Scotland e.g. NHS and self-management groups, which may help to reduce biased results towards a specific demographic. Although a larger sample recruited from the community to increase the generalisability of findings would have helped to increase reliability of the study.

The sample size was small due to a low response rate, however this is consistent with a postal survey response in a chronic illness group.⁵² The method for data collection may have created a sample bias where by individuals with greater functional disability were unable to

participate. The cross-sectional design prevents causal inference, therefore a suggestion for future research could conduct controlled studies in a laboratory setting where self-compassion could be manipulated and the effects on key outcomes systematically assessed (e.g., severity of pain, distress, quality of life, disability). In addition, the present study relied on self-report measures. Future studies investigating the relationship between self-compassion and pain adjustment should consider integrating self-report measures with direct measures, such as functional performance and structured clinical interviews.

The results of the current study indicate a need for further research particularly in relation to longitudinal and experimental studies, explicitly testing causal chains such as those proposed in this study. This was the first study to explore self-warmth, self-coldness and self-esteem as predictors in a chronic pain sample, replication with a larger sample is necessary to strengthen the conclusion drawn and to ensure generalisability.

2.4.4 Conclusion

The current study expands the literature by demonstrating self-warmth as a unique predictor of quality of life when distress, severity of pain and gender were controlled for in a chronic pain population. It was also found that neither, self-warmth, self-coldness or self-esteem were unique predictors of disability when controlling for distress and severity of pain. The findings suggest that interventions which may increase self-warmth such as Compassionate Mind Training ¹⁰¹ would be an effective treatment in addressing poor quality of life for people with chronic pain.

2.5 References

1. The Scottish Intercollegiate Guidelines Network (SIGN) Management of Chronic Pain. SIGN publication no136 [Internet] 136: 2013. Available from: <http://sign.ac.uk>
2. Breivik H, Collett B, Ventafridda V, Cohen R, Gallacher D: Survey of chronic pain in Europe: Prevalence, impact on daily life, and treatment. *European J of Pain* 2006; 10:287-287.
3. Wong W, Fielding R: Prevalence and characteristics of chronic pain in the general population of Hong Kong. *The J of Pain* 2011; 12:236-245.
4. Breivik H, Eisenberg E, O'Brien T: The individual and societal burden of chronic pain in Europe: The case for strategic prioritisation and action to improve knowledge and availability of appropriate care. *BMC Public Health* 2013; 13:1229.
5. Goldberg DMcGee S: Pain as a global public health priority. *BMC Public Health* 2011; 11:770.
6. Linton S, Nicholas M, MacDonald S: Development of a short form of the Örebro Musculoskeletal Pain Screening Questionnaire. *Spine* 2011; 36:1891-1895.
7. Jensen M, Turk D: Contributions of psychology to the understanding and treatment of people with chronic pain: Why it matters to all psychologists. *American Psychologist* 2014; 69:105-118.
8. Williams A, Eccleston C, Morley S: Systematic Review and meta-analysis of psychological treatments for persistent pain in adults, excluding headache. *Eur J of Pain* 2009; 13: S284.
9. McCracken L, Morley S: The Psychological Flexibility model: A basis for integration and progress in psychological approaches to chronic pain management. *The J of Pain* 2014; 15:221-234.
10. Yu L, Norton S, Harrison A, McCracken L: In search of the person in pain: A systematic review of conceptualization, assessment methods, and evidence for self and identity in chronic pain. *J of Contextual Beh Science* 2015; 4:246-262.
11. Fredrickson B: Cultivating positive emotions to optimize health and well-being. *Prevention & Treatment* 3: 2000.

12. Pinto-Gouveia J, Duarte C, Matos M, Fráguas S: The protective role of self-compassion in relation to psychopathology symptoms and quality of life in chronic and in cancer patients. *Clinical Psych & Psychother* 2013; 21:311-323.
13. Cho C, Seo H, Bae K, Lee K, Hwang I, Warner J. The impact of depression and anxiety on self-assessed pain, disability, and quality of life in patients scheduled for rotator cuff repair. *Journal of Shoulder and Elbow Surgery* 2013; 22:1160-1166.
14. Outcalt S, Kroenke K, Krebs E, Chumbler N, Wu J, Yu Z, Bair M. Chronic pain and comorbid mental health conditions: Independent associations of posttraumatic stress disorder and depression with pain, disability, and quality of life. *J Behav Med* 2015; 38:535-543.
15. Yarnell L, Stafford R, Neff K, Reilly E, Knox M, Mullarkey M: Meta-Analysis of gender differences in self-compassion. *Self and Identity* 2015; 14:499-520.
16. Neff, K, Pommier, E: The relationships between self-compassion and other focused concern among college undergraduates, community adults, and practicing meditators. *Self and Identity* 2013; 12:160-176.
17. Ardelt M: Are Older Adults Wiser than college students? A comparison of two age cohorts. *Journal of Adult Development* 2009; 17:193-207.
18. Neff K: Self-Compassion: An alternative conceptualization of a healthy attitude toward oneself. *Self and Identity* 2003; 2:85-101.
19. Brion J, Leary M, Drabkin A: Self-compassion and reactions to serious illness: The case of HIV. *J of Health Psychology* 2013; 19:218-229.
20. Diedrich A, Grant M, Hofmann S, Hiller W, Berking M: Self-compassion as an emotion regulation strategy in major depressive disorder. *Beh Res Ther* 2014; 58:43-51
21. MacBeth A, Gumley A: Exploring compassion: A meta-analysis of the association between self-compassion and psychopathology. *Clin Psych Rev* 2012; 32:545-552.
22. Neff K: The development and validation of a scale to measure self-compassion. *Self and Identity* 2003; 2:223-250.
23. Raes F, Pommier E, Neff K, Van Gucht D. Construction and factorial validation of a short form of the Self-Compassion Scale. *Clin Psych & Psychother* 2011; 18:250-255.

24. Muris P, Petrocchi N: Protection or Vulnerability? A Meta-Analysis of the relations between the positive and negative components of self-compassion and psychopathology. *Clin Psych & Psychother* (In Press).
25. Van Sonderen E, Sanderman R, Coyne JC. Ineffectiveness of reverse wording of questionnaire items: Let's learn from cows in the rain. *PloS one*. 2013;8(7): e68967.
26. López A, Sanderman R, Smink A, Zhang Y, van Sonderen E, Ranchor A, Schroevers M. A Reconsideration of the Self-Compassion Scale's total score: Self-Compassion versus Self-Criticism. *PLOS ONE* 10: e0132940, 2015.
27. Williams M, Dalgleish T, Karl A, Kuyken W: Examining the factor structures of the Five Facet Mindfulness Questionnaire and the Self-Compassion Scale. *Psych Assess* 2014; 26:407-418.
28. Klimecki O, Leiberg S, Lamm C, Singer T: Functional neural plasticity and associated changes in positive affect after compassion training. *Cerebral Cortex* 2012; 23:1552-1561.
29. Odou N, Brinker J: Exploring the relationship between rumination, self-compassion, and mood. *Self and Identity* 2014; 13:449-459.
30. Leaviss J, Uttley L: Psychotherapeutic benefits of Compassion-Focused Therapy: An early systematic review. *Psych Med* 2014; 45:927-945.
31. Terry M, Leary M: Self-compassion, self-regulation, and health. *Self and Identity* 2011; 10:352-362.
32. Doran N: Experiencing wellness within illness: Exploring a mindfulness-based approach to chronic back pain. *Qual Health Res* 2014; 24:749-760.
33. McLaren S, Gomez R, Bailey M, Van Der Horst R: The association of depression and sense of belonging with suicidal ideation among older adults: Applicability of resiliency models. *Suicide and Life-Threatening Behaviour* 2007; 37:89-102.
34. Castilho P, Pinto-Gouveia J, Duarte J: Evaluating the multifactor structure of the long and short versions of the Self-Compassion Scale in a clinical sample. *Journal of Clin Psych* 2015; 71:856-870.
35. Neff K, Vonk R: Self-compassion versus global self-esteem: Two different ways of relating to oneself. *Journal of Personality* 2009; 77:23-50.

36. Barry C, Doucette H, Loflin D, Rivera-Hudson N, Herrington L.: "Let Me Take a Selfie": Associations between self-photography, narcissism, and self-esteem. *Psychology of Popular Media Culture* 2015.
37. Marshall T, Lefringhausen K, Ferenczi N: The big five, self-esteem, and narcissism as predictors of the topics people write about in Facebook status updates. *Personality and Individual Differences* 2015; 85:35-40.
38. Juth V, Smyth J, Santuzzi A: How Do You Feel? Self-esteem predicts affect, stress, social interaction, and symptom severity during daily life in patients with chronic illness. *Journal of Health Psychology* 2008; 13:884-894.
39. Bisschop M, Kriegsman D, Beekman A, Deeg D: Chronic diseases and depression: the modifying role of psychosocial resources. *Social Science & Medicine* 2004; 59:721-733.
40. Simpson J, Lekwuwa G, Crawford T. Illness beliefs and psychological outcome in people with Parkinson's disease. *Chronic Illness* 2013; 9:165-176.
41. Chen WT, Wantland D, Reid P, Corless IB, Eller LS, Liping S, Holzemer WL, Nokes K, Sefcik E, Rivero-Mendez M, Voss J. Engagement with health care providers affects self-efficacy, self-esteem, medication adherence and quality of life in people living with HIV. *Journal of AIDS & Clinical Research*. 2013; 4:256.
42. Mikula P, Nagyova I, Krokavcova M, Vitkova M, Rosenberger J, Szilasiova J, Gdovinova Z, Stewart R, Groothoff J, van Dijk J. Self-esteem, social participation, and quality of life in patients with multiple sclerosis. *Journal of Health Psychology* (In Press).
43. Platten M, Newman E, Quayle E. Self-esteem and its relationship to mental health and quality of life in adults with Cystic Fibrosis. *J Clin Psychol Med Settings* 2012; 20:392-399.
44. Piquart M: Self-esteem of children and adolescents with chronic illness: a meta-analysis. *Child: Care, Health and Development* 2012; 39:153-161.
45. Bandura A: Editorial. *American Journal of Health Promotion* 1997; 12:8-10.
46. Hogg M, Vaughan G: MA Hogg & GM Vaughan (Eds.), *Social Psychology*. 6th ed. page 111-146, 2011.

47. Canetti L, Berry E, Elizur Y: Psychosocial predictors of weight loss and psychological adjustment following bariatric surgery and a weight-loss program: The mediating role of emotional eating. *Int J Eat Disorders* 2009; 42:109-117.
48. Neff KD. Self-compassion, self-esteem, and well-being. *Social and personality psychology compass*. 2011; 5:1-2.
49. Crocker J, Park L: The Costly Pursuit of Self-Esteem. *Psychological Bulletin* 2004; 130:392-414.
50. Baumeister R, Campbell J, Krueger J, Vohs K: Does High Self-Esteem Cause Better Performance, Interpersonal Success, Happiness, or Healthier Lifestyles? *Psychological Science in the Public Interest* 2003; 4:1-44.
51. Leary M, Tate E, Adams C, Batts Allen A, Hancock J. Self-compassion and reactions to unpleasant self-relevant events: The implications of treating oneself kindly. *Journal of Personality and Social Psychology* 2007; 92:887-904.
52. Hayter MDorstyn D. Resilience, self-esteem and self-compassion in adults with spina bifida. *Spinal Cord* 2013; 52:167-171.
53. Neff K, Pisitsungkagarn K, Hsieh Y: Self-compassion and self-construal in the United States, Thailand, and Taiwan. *Journal of Cross-Cultural Psychology* 2008; 39:267-285.
54. Attal N, Lanteri-Minet M, Laurent B, Fermanian J, Bouhassira D: The specific disease burden of neuropathic pain: Results of a French nationwide survey. *Pain* 2011; 152:2836-2843.
55. Hofmann S, Asnaani A, Vonk I, Sawyer A, Fang A. The Efficacy of Cognitive Behavioral Therapy: A review of meta-analyses. *Cognitive Therapy and Research* 2012; 36:427-440.
56. Falsarella G, Coimbra I, Neri A, Barcelos C, Costallat L, Carvalho O, Coimbra A. Impact of rheumatic diseases and chronic joint symptoms on quality of life in the elderly. *Archives of Gerontology and Geriatrics* 54: e77-e82, 2012.
57. Peterlin B, Tietjen G, Brandes J, Rubin S, Drexler E, Lidicker J, Meng S: Post traumatic stress disorder in migraine. *The Journal of Head and Face Pain* 2009; 49:541-551.
58. Burke A, Mathias J, Denson L: Psychological functioning of people living with chronic pain: A meta-analytic review. *Brit J of Clin Psych* 2015; 54:345-360.

59. Costa J, Marôco J, Pinto-Gouveia J, Ferreira C, Castilho P: Validation of the psychometric properties of the Self-Compassion Scale. Testing the factorial validity and factorial invariance of the measure among Borderline Personality Disorder, Anxiety Disorder, Eating Disorder and General Populations. *Clinical Psychology & Psychotherapy* 2015.
60. Wren A, Somers T, Wright M, Goetz M, Leary M, Fras A, Huh B, Rogers L, Keefe F. Self-compassion in patients with persistent musculoskeletal pain: Relationship of self-compassion to adjustment to persistent pain. *Journal of Pain and Symptom Management* 2012; 43:759-770.
61. Cleeland C, Ryan K. Pain assessment: Global use of the Brief Pain Inventory. *Annual Academic Medical Singapore* 1994; 23:129-138.
62. Wetherell J, Afari N, Rutledge T, Sorrell J, Stoddard J, Petkus A, Solomon B, Lehman D, Liu L, Lang A, Atkinson H. A randomized, controlled trial of acceptance and commitment therapy and cognitive-behavioral therapy for chronic pain. *Pain* 2011; 152:2098-2107.
63. Purdie F, Morley S: Self-compassion, pain, and breaking a social contract. *PAIN* 2015; 156:2354-2363.
64. Rosenberg M: *Society and the adolescent self-image*. Princeton, N.J.: Princeton University Press; 1965.
65. Sinclair, S. J., Blais, M. A., Gansler, D. A., Sandberg, E., Bistis, K., & LoCicero, A. (2010). Psychometric properties of the Rosenberg Self-Esteem Scale: Overall and across demographic groups living within the United States. *Evaluation and the Health Professions*, 33, 56–80.
66. Eller L, Rivero-Mendez M, Voss J, Chen W, Chaiphibalsarisdi P, Iipinge S. Depressive symptoms, self-esteem, HIV symptom management self-efficacy and self-compassion in people living with HIV. *AIDS Care* 26:795-803, 2013.
67. Dworkin R, Turk D, Farrar J, Haythornthwaite J, Jensen M. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain* 2005; 113:9-19.
68. Tan G, Jensen M, Thornby J, Shanti B. Validation of the brief pain inventory for chronic nonmalignant pain. *The Journal of Pain* 2004; 5:133-137.

69. Keller S, Bann C, Dodd S, Schein J, Mendoza T, Cleeland C: Validity of the Brief Pain Inventory for use in documenting the outcomes of patients with non-cancer pain. *The Clinical Journal of Pain* 2004; 20:309-318.
70. Zigmond Asnaith R. The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica* 1983; 67:361-370.
71. Clover K, Carter G, Mackinnon A, Adams C: Is my patient suffering clinically significant emotional distress? Demonstration of a probabilities approach to evaluating algorithms for screening for distress. *Support Care Cancer* 2009; 17:1455-1462.
72. Turk D, Dworkin R, Trudeau J, Benson C, Biondi D, Katz N, Kim M. Validation of the Hospital Anxiety and Depression Scale in patients with acute low back pain. *The Journal of Pain* 2015; 16:1012-1021.
73. Gillanders D, Sinclair A, MacLean M, Jardine K. Illness cognitions, cognitive fusion, avoidance and self-compassion as predictors of distress and quality of life in a heterogeneous sample of adults, after cancer. *Journal of Contextual Behavioral Science* 2015; 4:300-311.
74. Anagnostis C, Gatchel R, Mayer T: The Pain Disability Questionnaire. *Spine* 2004; 29:2290-2302.
75. Gatchel R, Mayer T, Theodore B. The Pain Disability Questionnaire: Relationship to one-year functional and psychosocial rehabilitation outcomes. *Journal of Occupational Rehabilitation* 2006; 16:72-91.
76. Burckhardt C, Anderson K, Archenholtz B, Hägg O: Health and quality of life outcomes 2003; 1:59.
77. Giordano PC, Alexandre NM, Rodrigues RC, Coluci MZ. The Pain Disability Questionnaire: a reliability and validity study. *Revista Latino-Americana de Enfermagem*. 2012; 20:76-83.
78. Bentsen S, Miaskowski C, Rustøen T: Demographic and clinical characteristics associated with quality of life in patients with chronic obstructive pulmonary disease. *Qual Life Res* 2013; 23:991-998.

79. Liedberg G, Burckhardt C, Henriksson C: Validity and reliability testing of the Quality of Life Scale, Swedish version in women with fibromyalgia - statistical analyses. *Scandinavian Journal of Caring Sciences* 2005; 19:64-70.
80. Hayes A. *Introduction to Mediation, Moderation, and Conditional Process Analysis*, 2013.
81. Preacher K, Hayes A. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods* 2008; 40:879-891.
82. Shrive F, Stuart H, Quan H, Ghali W. Dealing with missing data in a multi-question depression scale: A comparison of imputation methods. *BMC Medical Research Methodology* 2015; 6:57.
83. Field A. *Discovering Statistics Using IBM SPSS Statistics*, 2013.
84. Cairns BGazerani P: Sex-related differences in pain. *Maturitas* 2009; 63:292-296.
85. García-Martínez A, De Paz J, Márquez S. Effects of an exercise programme on self-esteem, self-concept and quality of life in women with fibromyalgia: a randomized controlled trial. *Rheumatol Int* 2011; 32:1869-1876.
86. Kazemi H, Ghasemi S, Fereshtehnejad S: Anxiety and depression in patients with amputated limbs suffering from phantom pain: A comparative study with non-phantom chronic pain. *International Journal of Preventive Medicine* 2013; 4:218-225.
87. Börsbo B, Gerdle B, Peolsson M. Impact of the interaction between self-efficacy, symptoms and catastrophising on disability, quality of life and health in with chronic pain patients. *Disability and Rehabilitation* 2010; 32:1387-1396.
88. Goesling J, Clauw D, Hassett A: Pain and depression: An integrative review of neurobiological and psychological factors. *Current Psychiatry Reports* 15: 2013.
89. Skevington S, McCrate F. Expecting a good quality of life in health: Assessing people with diverse diseases and conditions using the WHOQOL-BREF. *Health Expectations* 2011; 15:49-62.
90. Rustøen T, Wahl A, Hanestad B, Lerdal A, Paul S, Miaskowski C. Gender differences in chronic pain—findings from a population-based study of Norwegian adults. *Pain Management Nursing* 2004; 5:105-117.

91. Unruh A, Ritchie J, Merskey H: Does gender affect appraisal of pain and pain coping strategies? *The Clinical Journal of Pain* 15:31-40, 1999.
92. Fillingim R, King C, Ribeiro-Dasilva M, Rahim-Williams B, Riley J: Sex, gender, and pain: A review of recent clinical and experimental findings. *The Journal of Pain* 2009; 10:447-485.
93. Muris P, Meesters C, Pierik A, de Kock B: Good for the self: Self-Compassion and other self-related constructs in relation to symptoms of anxiety and depression in non-clinical youths. *Journal of Child and Family Studies* 2015; 25:607-617.
94. Hwang S, Kim G, Yang J, Yang E: The moderating effects of age on the relationships of self-compassion, self-esteem, and mental health. *Japanese Psychological Research* 2016; 58:194-205.
95. Leary M, Tangney J: *Handbook of self and identity*. 2nd ed. New York: Guilford Press; 2012.
96. van den Beuken-van Everdingen M, de Rijke J, Kessels A, Schouten H, van Kleef M, Patijn J. Prevalence of pain in patients with cancer: a systematic review of the past 40 years. *Annals of Oncology* 2007; 18:1437-1449.
97. Ware J, Kosinski M, Dewey J. How to score version 2 of the SF-36 health survey (standard & acute forms). Lincoln, R.I.: QualityMetric Inc.; 2000.
98. Bartoces M, Severson R, Rusin B, Schwartz K, Ruterbusch J, Neale A. Quality of life and self-esteem of long-term survivors of invasive and non-invasive cervical cancer. *Journal of Women's Health* 2009; 18:655-661.
99. Hilbert A, Braehler E, Schmidt R, Lowe B, Hauser W, Zenger M. Self-Compassion as a resource in the self-stigma process of overweight and obese individuals. *Obesity Facts* 2015; 8:293-301.
100. Hoge E, Hölzel B, Marques L, Metcalf C, Brach N, Lazar S, Simon N. Mindfulness and self-compassion in Generalized Anxiety Disorder: Examining predictors of disability. *Evidence-Based Complementary and Alternative Medicine* 2013; 1-7.
101. Gilbert Procter S: Compassionate mind training for people with high shame and self-criticism: overview and pilot study of a group therapy approach. *Clin Psych & Psychother* 2006; 13:353-379.

102. Carson JW, Keefe FJ, Lynch TR, Carson KM, Goli V, Fras AM, Thorp SR. Loving-kindness meditation for chronic low back pain results from a pilot trial. *Journal of Holistic Nursing*. 2005; 23:287-304.
103. Chapin HL, Darnall BD, Seppala EM, Doty JR, Hah JM, Mackey SC. Pilot study of a compassion meditation intervention in chronic pain. *Journal of Compassionate Health Care*. 2014; 1:1.
104. Flink I, Smeets E, Bergbom S, Peters M. Happy despite pain: Pilot study of a positive psychology intervention for patients with chronic pain. *Scandinavian Journal of Pain* 2015; 7:71-79.
105. Linton S: Applying dialectical behavior therapy to chronic pain: A case study. *Scandinavian Journal of Pain* 2010; 1:50-54.
106. McCracken LM, Sato A, Taylor GJ. A trial of a brief group-based form of Acceptance and Commitment Therapy (ACT) for chronic pain in general practice: Pilot outcome and process results. *J of Pain* 2013; 14:1398-406.
107. Barnard L, Curry J: Self-compassion: Conceptualizations, correlates, & interventions. *Review of General Psychology* 2011; 15:289-303.
108. Nowell R. SPICE briefing "Mental Health in Scotland" [Internet]. Scottish Parliament Website. 2014 [cited 1 March 2016]. Available from: http://www.scottish.parliament.uk/ResearchBriefingsAndFactsheets/S4/SB_14-36.pdf

Full Thesis Reference List

- Ablin, J., Fitzcharles, M., Buskila, D., Shir, Y., Sommer, C., & Häuser, W. (2013). Treatment of Fibromyalgia Syndrome: Recommendations of Recent Evidence-Based Interdisciplinary Guidelines with Special Emphasis on Complementary and Alternative Therapies. *Evidence-Based Complementary And Alternative Medicine*, 2013, 1-7. <http://dx.doi.org/10.1155/2013/485272>
- Akkaya, N., Akkaya, S., Atalay, N., Balci, C., & Sahin, F. (2012). Relationship between the body image and level of pain, functional status, severity of depression, and quality of life in patients with fibromyalgia syndrome. *Clin Rheumatol*, 31(6), 983-988.
- Alanoğlu, E., Ulaş, U., Özdağ, F., Odabaşı, Z., Çakıcı, A., & Vural, O. (2004). Auditory event-related brain potentials in fibromyalgia syndrome. *Rheumatol Int*, 25(5), 345-349.
- Alonso, J., Ferrer, M., Gandek, B., Ware Jr., J., Aaronson, N., & Mosconi, P. et al. (2004). Health-related quality of life associated with chronic conditions in eight countries: Results from the International Quality of Life Assessment (IQOLA) Project. *Qual Life Res*, 13(2), 283-298.
- Anagnostis, C., Gatchel, R., & Mayer, T. (2004). The Pain Disability Questionnaire. *Spine*, 29(20), 2290-2302.
- Andrell, P., Schultz, T., Mannerkorpi, K., Nordeman, L., Borjesson, M., & Mannheimer, C. (2014). Health-related quality of life in fibromyalgia and refractory angina pectoris: A comparison between two chronic non-malignant pain disorders. *Journal of Rehabilitation Medicine*, 46(4), 341-347.
- Ardelt, M. (2009). Are Older Adults Wiser Than College Students? A Comparison of Two Age Cohorts. *Journal of Adult Development*, 17(4), 193-207.
- Attal, N., Lanteri-Minet, M., Laurent, B., Fermanian, J., & Bouhassira, D. (2011). The specific disease burden of neuropathic pain: Results of a French nationwide survey. *Pain*, 152(12), 2836-2843.
- Avila, L., de Araujo Filho, G., Guimarães, E., Gonçalves, L., Paschoalin, P., & Aleixo, F. (2014). Characterization of the pain, sleep and alexithymia patterns of patients with fibromyalgia treated in a Brazilian tertiary center. *Revista Brasileira De Reumatologia (English Edition)*, 54, 409-413.
- Bandura, A. (1997). Editorial. *American Journal of Health Promotion*, 12, 8-10.

- Barnard, L. & Curry, J. (2011). Self-compassion: Conceptualizations, correlates, & interventions. *Review of General Psychology, 15*, 289-303.
- Barry, C., Doucette, H., Loflin, D., Rivera-Hudson, N., & Herrington, L. (2015). "Let me take a selfie": Associations between self-photography, narcissism, and self-esteem. *Psychology of Popular Media Culture*.
- Bartoces, M., Severson, R., Rusin, B., Schwartz, K., Ruterbusch, J., & Neale, A. (2009). Quality of life and self-esteem of long-term survivors of invasive and non-invasive cervical cancer. *Journal of Women's Health, 18*, 655-661.
- Batmaz, İ., Saryıldız, M., Dilek, B., İnanır, A., Demircan, Z., & Hatipoğlu, N. et al. (2012). Sexuality of men with fibromyalgia: What are the factors that cause sexual dysfunction? *Rheumatology International, 33*, 1265-1270.
- Baumeister, R., Campbell, J., Krueger, J., & Vohs, K. (2003). Does high self-esteem cause better performance, interpersonal success, happiness, or healthier lifestyles? *Psychological Science in The Public Interest, 4*, 1-44.
- Bennett, R., Friend, R., Jones, K., Ward, R., Han, B., & Ross, R. (2009). The Revised Fibromyalgia Impact Questionnaire (FIQR): validation and psychometric properties. *Arthritis Research Therapy, 11*, 415.
- Bentsen, S., Miaskowski, C., & Rustøen, T. (2013). Demographic and clinical characteristics associated with quality of life in patients with chronic obstructive pulmonary disease. *Quality of Life Research, 23*, 991-998.
- Berzon, R., Hays, R., & Shumaker, S. (1993). International use, application and performance of health-related quality of life instruments. *Quality of Life Research, 2*, 367-368.
- Birtane, M., Uzunca, K., Taştekin, N., & Tuna, H. (2006). The evaluation of quality of life in fibromyalgia syndrome: a comparison with rheumatoid arthritis by using SF-36 Health Survey. *Clinical Rheumatology, 26*, 679-684.
- Bisschop, M., Kriegsman, D., Beekman, A., & Deeg, D. (2004). Chronic diseases and depression: the modifying role of psychosocial resources. *Social Science & Medicine, 59*, 721-733. <http://dx.doi.org/10.1016/j.socscimed.2003.11.038>
- Börsbo, B., Gerdle, B., & Peolsson, M. (2010). Impact of the interaction between self-efficacy, symptoms and catastrophising on disability, quality of life and health in with chronic pain patients. *Disability and Rehabilitation, 32*, 1387-1396.

- Branco, J., Bannwarth, B., Failde, I., Abello Carbonell, J., Blotman, F., & Spaeth, M. et al. (2010). Prevalence of Fibromyalgia: A Survey in Five European Countries. *Seminars in Arthritis and Rheumatism*, 39, 448-453.
- Breivik, H., Collett, B., Ventafridda, V., Cohen, R., & Gallacher, D. (2006). Survey of chronic pain in Europe: Prevalence, impact on daily life, and treatment. *European Journal of Pain*, 10, 287-287.
- Breivik, H., Eisenberg, E., & O'Brien, T. (2013). The individual and societal burden of chronic pain in Europe: the case for strategic prioritisation and action to improve knowledge and availability of appropriate care. *BMC Public Health*, 13, 1229.
- Brion, J., Leary, M., & Drabkin, A. (2013). Self-compassion and reactions to serious illness: The case of HIV. *Journal of Health Psychology*, 19, 218-229.
- Bullinger, M., Alonso, J., Apolone, G., Leplège, A., Sullivan, M., & Wood-Dauphinee, S. et al. (1998). Translating health status questionnaires and evaluating their quality. *Journal of Clinical Epidemiology*, 51, 913-923.
- Burckhardt, C., Anderson, K., Archenholtz, B., & Hägg, O. (2003). *Health and Quality of Life Outcomes*, 1, 59.
- Burckhardt, C., Woods, S., Schultz, A., & Ziebarth, D. (1989). Quality of life of adults with chronic illness: A psychometric study. *Research Nursing Health*, 12, 347-354.
- Burke, A., Mathias, J., & Denson, L. (2015). Psychological functioning of people living with chronic pain: A meta-analytic review. *British Journal of Clinical Psychology*, 54, 345-360.
- Cairns, B. & Gazerani, P. (2009). Sex-related differences in pain. *Maturitas*, 63, 292-296.
- Calandre, E., Garcia-Carrillo, J., Garcia-Leiva, J., Rico-Villademoros, F., Molina-Barea, R., & Rodriguez-Lopez, C. (2010). Subgrouping patients with fibromyalgia according to the results of the fibromyalgia impact questionnaire: a replication study. *Rheumatology International*, 31, 1555-1559.
- Campos, R. & Vázquez, M. (2011). Health-related quality of life in women with fibromyalgia: clinical and psychological factors associated. *Clinical Rheumatology*, 31, 347-355.
- Canetti, L., Berry, E., & Elizur, Y. (2009). Psychosocial predictors of weight loss and psychological adjustment following bariatric surgery and a weight-loss program: The mediating role of emotional eating. *International Journal of Eating Disorders*, 42, 109-117.

- Cardoso, F., Curtolo, M., Natour, J., & Lombardi Júnior, I. (2011). Avaliação da qualidade de vida, força muscular e capacidade funcional em mulheres com fibromialgia. *Rev. Bras. Reumatol.*, 51, 344-350.
- Carrillo-de-la-Peña, M., Triñanes, Y., González-Villar, A., Romero-Yuste, S., Gómez-Perretta, C., Arias, M., & Wolfe, F. (2014). Convergence between the 1990 and 2010 ACR diagnostic criteria and validation of the Spanish version of the Fibromyalgia Survey Questionnaire (FSQ). *Rheumatology International*, 35, 141-151.
- Castilho, P., Pinto-Gouveia, J., & Duarte, J. (2015). Evaluating the Multifactor Structure of the Long and Short Versions of the Self-Compassion Scale in a Clinical Sample. *Journal Of Clinical Psychology*, 71(9), 856-870
- Cherepanov, D., Palta, M., Fryback, D., Robert, S., Hays, R., & Kaplan, R. (2011). Gender Differences in Multiple Underlying Dimensions of Health-related Quality of Life Are Associated with Sociodemographic and Socioeconomic Status. *Medical Care*, 49(11), 1021-1030.
- Cho, C., Seo, H., Bae, K., Lee, K., Hwang, I., & Warner, J. (2013). The impact of depression and anxiety on self-assessed pain, disability, and quality of life in patients scheduled for rotator cuff repair. *Journal Of Shoulder And Elbow Surgery*, 22(9), 1160-1166.
- Clauw, D. (2014). Fibromyalgia. *JAMA*, 311(15), 1547. <http://dx.doi.org/10.1001/jama.2014.3266>
- Cleeland, C. & Ryan, K. (1994). Pain assessment: Global use of the Brief Pain Inventory. *Annual Academic Medical Singapore*, 23(2), 129-138.
- Clover, K., Carter, G., Mackinnon, A., & Adams, C. (2009). Is my patient suffering clinically significant emotional distress? Demonstration of a probabilities approach to evaluating algorithms for screening for distress. *Support Care Cancer*, 17(12), 1455-1462.
- Costa, J., Marôco, J., Pinto-Gouveia, J., Ferreira, C., & Castilho, P. (2015). Validation of the Psychometric Properties of the Self-Compassion Scale. Testing the Factorial Validity and Factorial Invariance of the Measure among Borderline Personality Disorder, Anxiety Disorder, Eating Disorder and General Populations. *Clinical Psychology & Psychotherapy*,
- Crocker, J. & Park, L. (2004). The Costly Pursuit of Self-Esteem. *Psychological Bulletin*, 130(3), 392-
- de Souza, J., Goffaux, P., Julien, N., Potvin, S., Charest, J., & Marchand, S. (2008). Fibromyalgia subgroups: profiling distinct subgroups using the Fibromyalgia Impact Questionnaire. A

- preliminary study. *Rheumatol Int*, 29(5), 509-515. <http://dx.doi.org/10.1007/s00296-008-0722-5>
- Dean Wantland, W. (2013). Engagement with Health Care Providers Affects Self- Efficacy, Self-Esteem, Medication Adherence and Quality of Life in People Living with HIV. *J AIDS Clin Res*, 04(01)
- DerSimonian, R. & Laird, N. (1986). Meta-analysis in clinical trials. *Controlled Clinical Trials*, 7(3), 177-188.
- Diedrich, A., Grant, M., Hofmann, S., Hiller, W., & Berking, M. (2014). Self-compassion as an emotion regulation strategy in major depressive disorder. *Behaviour Research And Therapy*, 58, 43-51.
- Doran, N. (2014). Experiencing Wellness Within Illness: Exploring a Mindfulness-Based Approach to Chronic Back Pain. *Qualitative Health Research*, 24(6), 749-760.
- Dworkin, R., Turk, D., Farrar, J., Haythornthwaite, J., Jensen, M., & Katz, N. et al. (2005). Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain*, 113(1), 9-19.
- Egger, M., Smith, G.D., Schneider, M., Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315, 629-634.
- Eller, L., Rivero-Mendez, M., Voss, J., Chen, W., Chaiphibalsarisdi, P., & Ipinge, S. et al. (2013). Depressive symptoms, self-esteem, HIV symptom management self-efficacy and self-compassion in people living with HIV. *AIDS Care*, 26(7), 795-803.
- Elm, E., Altman, D., Egger, M., Pocock, S., Gotsche, P., & Vandenbroucke, J. (2007). Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*, 335(7624), 806-808.
- Falsarella, G., Coimbra, I., Neri, A., Barcelos, C., Costallat, L., Carvalho, O., & Coimbra, A. (2012). Impact of rheumatic diseases and chronic joint symptoms on quality of life in the elderly. *Archives Of Gerontology And Geriatrics*, 54(2), e77-e82.
- Faust, A., Halpern, L., Danoff-Burg, S., & Cross, R. (2012). Psychosocial factors contributing to inflammatory bowel disease activity and health-related quality of life. *Gastroenterology & Hepatology*, 8(3), 173-181. Retrieved from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3365520/>

Field, A. *Discovering statistics using IBM SPSS statistics*.

Fillingim, R., King, C., Ribeiro-Dasilva, M., Rahim-Williams, B., & Riley, J. (2009). Sex, Gender, and Pain: A Review of Recent Clinical and Experimental Findings. *The Journal Of Pain*, 10(5), 447-485. <http://dx.doi.org/10.1016/j.jpain.2008.12.001>

Flink, I., Smeets, E., Bergbom, S., & Peters, M. (2015). Happy despite pain: Pilot study of a positive psychology intervention for patients with chronic pain. *Scandinavian Journal Of Pain*, 7, 71-79. <http://dx.doi.org/10.1016/j.sjpain.2015.01.005>

Fredrickson, B. (2000). Cultivating positive emotions to optimize health and well-being. *Prevention & Treatment*, 3(1). <http://dx.doi.org/10.1037/1522-3736.3.1.31a>

Fuller-Thomson, E., Nimigon-Young, J., & Brennenstuhl, S. (2011). Individuals with fibromyalgia and depression: Findings from a nationally representative Canadian survey. *Rheumatol Int*, 32(4), 853-862.

García-Martínez, A., De Paz, J., & Márquez, S. (2011). Effects of an exercise programme on self-esteem, self-concept and quality of life in women with fibromyalgia: a randomized controlled trial. *Rheumatol Int*, 32(7), 1869-1876.

Garratt, A., Ruta, D., Abdalla, M., Buckingham, J., & Russell, I. (1993). The SF36 health survey questionnaire: an outcome measure suitable for routine use within the NHS?. *BMJ*, 306(6890), 1440-1444.

Gatchel, R., Mayer, T., & Theodore, B. (2006). The Pain Disability Questionnaire: Relationship to One-Year Functional and Psychosocial Rehabilitation Outcomes. *Journal Of Occupational Rehabilitation*, 16(1), 72-91.

Gilbert, P. & Procter, S. (2006). Compassionate mind training for people with high shame and self-criticism: overview and pilot study of a group therapy approach. *Clinical Psychology & Psychotherapy*, 13(6), 353-379.

Gillanders, D., Sinclair, A., MacLean, M., & Jardine, K. (2015). Illness cognitions, cognitive fusion, avoidance and self-compassion as predictors of distress and quality of life in a heterogeneous sample of adults, after cancer. *Journal Of Contextual Behavioral Science*, 4(4), 300-311.

Goesling, J., Clauw, D., & Hassett, A. (2013). Pain and Depression: An Integrative Review of Neurobiological and Psychological Factors. *Current Psychiatry Reports*, 15(12). <http://dx.doi.org/10.1007/s11920-013-0421-0>

- Goldberg, D. & McGee, S. (2011). Pain as a global public health priority. *BMC Public Health*, 11(1), 770.
- Goldenberg, D. (2009). Diagnosis and Differential Diagnosis of Fibromyalgia. *The American Journal Of Medicine*, 122(12), S14-S21. <http://dx.doi.org/10.1016/j.amjmed.2009.09.007>
- Hayes, A. & Krippendorff, K. (2007). Answering the Call for a Standard Reliability Measure for Coding Data. *Communication Methods And Measures*, 1(1), 77-89.
- Hayes, A. (2013) *Introduction to mediation, moderation, and conditional process analysis*.
- Hayter, M. & Dorstyn, D. (2013). Resilience, self-esteem and self-compassion in adults with spina bifida. *Spinal Cord*, 52(2), 167-171.
- Higgins, J. (2003). Measuring inconsistency in meta-analyses. *BMJ*, 327(7414), 557-560.
- Hilbert, A., Braehler, E., Schmidt, R., Lowe, B., Hauser, W., & Zenger, M. (2015). Self-Compassion as a Resource in the Self-Stigma Process of Overweight and Obese Individuals. *Obesity Facts*, 8(5), 293-301.
- Hoffman, D. & Dukes, E. (2007). The health status burden of people with fibromyalgia: a review of studies that assessed health status with the SF-36 or the SF-12. *International Journal Of Clinical Practice*, 62(1), 115-126.
- Hofmann, S., Asnaani, A., Vonk, I., Sawyer, A., & Fang, A. (2012). The Efficacy of Cognitive Behavioral Therapy: A Review of Meta-analyses. *Cognitive Therapy And Research*, 36(5), 427-440.
- Hoge, E., Hölzel, B., Marques, L., Metcalf, C., Brach, N., Lazar, S., & Simon, N. (2013). Mindfulness and Self-Compassion in Generalized Anxiety Disorder: Examining Predictors of Disability. *Evidence-Based Complementary And Alternative Medicine*, 2013, 1-7.
- Hogg, M. & Vaughan, G. (2011). *MA Hogg & GM Vaughan (Eds.), Social Psychology* (6th ed., pp. 111-146).
- Hwang, S., Kim, G., Yang, J., & Yang, E. (2016). The Moderating Effects of Age on the Relationships of Self-Compassion, Self-Esteem, and Mental Health. *Japanese Psychological Research*, 58(2), 194-205.
- Jarde, A., Losilla, J., Vives, J., & F. Rodrigo, M. (2013). Q-Coh: A tool to screen the methodological quality of cohort studies in systematic reviews and meta-analyses. *International Journal Of Clinical And Health Psychology*, 13(2), 138-146.

- Jensen, M. & Turk, D. (2014). Contributions of psychology to the understanding and treatment of people with chronic pain: Why it matters to ALL psychologists. *American Psychologist*, 69(2), 105-118.
- Jüni, P., Witschi, A., Bloch, R., & Egger, M. (1999). The Hazards of Scoring the Quality of Clinical Trials for Meta-analysis. *JAMA*, 282(11), 1054.
- Juth, V., Smyth, J., & Santuzzi, A. (2008). How Do You Feel?: Self-esteem Predicts Affect, Stress, Social Interaction, and Symptom Severity during Daily Life in Patients with Chronic Illness. *Journal Of Health Psychology*, 13(7), 884-894.
- Kazemi, H., Ghasemi, S., & Fereshtehnejad, S. (2013). Anxiety and depression in patients with amputated limbs suffering from phantom pain: A comparative study with non-phantom chronic pain. *International Journal Of Preventive Medicine*, 4(2), 218-225.
- Keller, S., Bann, C., Dodd, S., Schein, J., Mendoza, T., & Cleeland, C. (2004). Validity of the Brief Pain Inventory for Use in Documenting the Outcomes of Patients With Noncancer Pain. *The Clinical Journal Of Pain*, 20(5), 309-318.
- Kim, S., Kim, S., Lee, C., Lee, H., Lee, S., & Park, Y. et al. (2012). Effect of fibromyalgia syndrome on the health-related quality of life and economic burden in Korea. *Rheumatology*, 52(2), 311-320.
- Klimecki, O., Leiberg, S., Lamm, C., & Singer, T. (2012). Functional Neural Plasticity and Associated Changes in Positive Affect After Compassion Training. *Cerebral Cortex*, 23(7), 1552-1561.
- Kmet, L., Lee, R., & Cook, L. (2004). *Institute of Health Economics. ihe.ca*. Retrieved 1 March 2016, from <http://www.ihe.ca/advanced-search/standard-quality-assessment-criteria-for-evaluating-primary-research-papers-from-a-variety-of-fields>
- Knudsen, A., Hotopf, M., Skogen, J., Overland, S., & Mykletun, A. (2010). The Health Status of Nonparticipants in a Population-based Health Study: The Hordaland Health Study. *American Journal Of Epidemiology*, 172(11), 1306-1314. <http://dx.doi.org/10.1093/aje/kwq257>
- Kohali, S., Khbazi, A., Hajaliloo, M., Namvar, L., & Farzin, H. (2010). The Evaluation Of Quality Of Life In Women With Rheumatoid Arthritis, Osteoarthritis And Fibromyalgia As Compared With Quality Of Life In Normal Women. *IJRH*, 7(1), 1-5. <http://dx.doi.org/10.5580/1ee9>
- Kucukali Turkyilmaz, A., Kurt, E., Karkucak, M., & Capkin, E. (2012). Sociodemographic Characteristics, Clinical Signs and Quality of Life in Patients with Fibromyalgia. *The Eurasian Journal Of Medicine*, 44(2), 88-93.

- Latorre-Román, P., Martínez-Amat, A., Martínez-López, E., Moral, Á., Santos, M., & Hita-Contreras, F. (2013). Validation and psychometric properties of the Spanish version of the Quality of Life Scale (QOLS) in patients with fibromyalgia. *Rheumatol Int*, 34(4), 543-549.
- Leary, M. & Tangney, J. (2012). *Handbook of self and identity* (2nd ed.). New York: Guilford Press.
- Leary, M., Tate, E., Adams, C., Batts Allen, A., & Hancock, J. (2007). Self-compassion and reactions to unpleasant self-relevant events: The implications of treating oneself kindly. *Journal Of Personality And Social Psychology*, 92(5), 887-904.
- Leaviss, J. & Uttley, L. (2014). Psychotherapeutic benefits of compassion-focused therapy: an early systematic review. *Psychological Medicine*, 45(05), 927-945.
- Liedberg, G., Burckhardt, C., & Henriksson, C. (2005). Validity and reliability testing of the Quality of Life Scale, Swedish version in women with fibromyalgia - statistical analyses. *Scandinavian Journal Of Caring Sciences*, 19(1), 64-70.
- Linton, S. (2010). Applying dialectical behavior therapy to chronic pain: A case study. *Scandinavian Journal Of Pain*, 1(1), 50-54.
- Linton, S., Nicholas, M., & MacDonald, S. (2011). Development of a Short Form of the Örebro Musculoskeletal Pain Screening Questionnaire. *Spine*, 36(22), 1891-1895.
- Linton, S., Nicholas, M., MacDonald, S., Boersma, K., Bergbom, S., Maher, C., & Refshaug, K. (2011). The role of depression and catastrophizing in musculoskeletal pain. *European Journal Of Pain*, 15(4), 416-422.
- López, A., Sanderman, R., Smerk, A., Zhang, Y., van Sonderen, E., Ranchor, A., & Schroevers, M. (2015). A Reconsideration of the Self-Compassion Scale's Total Score: Self-Compassion versus Self-Criticism. *PLOS ONE*, 10(7), e0132940.
- MacBeth, A. & Gumley, A. (2012). Exploring compassion: A meta-analysis of the association between self-compassion and psychopathology. *Clinical Psychology Review*, 32(6), 545-552.
- Mallen, C., Peat, G., & Croft, P. (2006). Quality assessment of observational studies is not commonplace in systematic reviews. *Journal Of Clinical Epidemiology*, 59(8), 765-769.
- Marshall, T., Lefringhausen, K., & Ferenczi, N. (2015). The Big Five, self-esteem, and narcissism as predictors of the topics people write about in Facebook status updates. *Personality And Individual Differences*, 85, 35-40.

- Mas, A., Carmona, L., Valverde, M., & Ribas, B. (2008). Prevalence and impact of fibromyalgia on function and quality of life in individuals from the general population: results from a nationwide study in Spain. *Clinical And Experimental Rheumatology*, 26(4), 519-526. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18799079>
- Matcham, F., Rayner, L., Hutton, J., Monk, A., Steel, C., & Hotopf, M. (2014). Self-help interventions for symptoms of depression, anxiety and psychological distress in patients with physical illnesses: A systematic review and meta-analysis. *Clinical Psychology Review*, 34(2), 141-157.
- McBeth, J. & Jones, K. (2006). Epidemiology of chronic musculoskeletal pain. *Best Practice & Research Clinical Rheumatology*, 21, 403-425.
- McCracken, L. & Morley, S. (2014). The Psychological Flexibility Model: A Basis for Integration and Progress in Psychological Approaches to Chronic Pain Management. *The Journal Of Pain*, 15(3), 221-234. <http://dx.doi.org/10.1016/j.jpain.2013.10.014>
- McLaren, S., Gomez, R., Bailey, M., & Van Der Horst, R. (2007). The Association of Depression and Sense of Belonging with Suicidal Ideation Among Older Adults: Applicability of Resiliency Models. *Suicide And Life-Threatening Behavior*, 37(1), 89-102.
- Mikula, P., Nagyova, I., Krokavcova, M., Vitkova, M., Rosenberger, J., & Szilasiova, J. et al. (2016). Self-esteem, social participation, and quality of life in patients with multiple sclerosis. *Journal Of Health Psychology*. <http://dx.doi.org/10.1177/1359105315621778>
- Moradkhani, A., Beckman, L., & Tabibian, J. (2013). Health-related quality of life in inflammatory bowel disease: Psychosocial, clinical, socioeconomic, and demographic predictors. *Journal Of Crohn's And Colitis*, 7(6), 467-473.
- Mottaghi, P., Qolipour, K., Safiri, S., Bahman Ziari, N., Kordi, A., & Izadi, S. et al. (2015). Quality of Life and Its Predictors Among Iranian Patients With Rheumatoid Arthritis: A Systematic Review. *Jentashapir Journal Of Health Research*, 6(2).
- Muris, P. & Petrocchi, N. (2016). Protection or Vulnerability? A Meta-Analysis of the Relations Between the Positive and Negative Components of Self-Compassion and Psychopathology. *Clinical Psychology & Psychotherapy*, (In Press) <http://dx.doi.org/10.1002/cpp.2005>

- Muris, P., Meesters, C., Pierik, A., & de Kock, B. (2015). Good for the Self: Self-Compassion and Other Self-Related Constructs in Relation to Symptoms of Anxiety and Depression in Non-clinical Youths. *Journal Of Child And Family Studies*, 25(2), 607-617.
- NEFF, K. (2003). Self-Compassion: An Alternative Conceptualization of a Healthy Attitude Toward Oneself. *Self And Identity*, 2(2), 85-101.
- NEFF, K. (2003). The Development and Validation of a Scale to Measure Self-Compassion. *Self And Identity*, 2(3), 223-250.
- Neff, K. & Vonk, R. (2009). Self-Compassion Versus Global Self-Esteem: Two Different Ways of Relating to Oneself. *Journal Of Personality*, 77(1), 23-50.
- Neff, K., Pisitsungkagarn, K., & Hsieh, Y. (2008). Self-Compassion and Self-Construal in the United States, Thailand, and Taiwan. *Journal Of Cross-Cultural Psychology*, 39(3), 267-285.
- Neumann, L., Berzak, A., & Buskila, D. (2000). Measuring health status in Israeli patients with fibromyalgiasyndrome and widespread pain and healthy individuals: Utility of the Short Form 36-item health survey (SF-36). *Seminars In Arthritis And Rheumatism*, 29(6), 400-408.
- Nowell, R. (2014). *SPICe briefing "Mental Health in Scotland"*. *Scottish Parliament Website*. Retrieved 1 March 2016, from http://www.scottish.parliament.uk/ResearchBriefingsAndFactsheets/S4/SB_14-36.pdf
- O'brien, R. (2007). A Caution Regarding Rules of Thumb for Variance Inflation Factors. *Quality & Quantity*, 41(5), 673-690.
- Odou, N. & Brinker, J. (2014). Exploring the Relationship between Rumination, Self-compassion, and Mood. *Self And Identity*, 13(4), 449-459.
- Okifuji, A., Bradshaw, D., Donaldson, G., & Turk, D. (2011). Sequential Analyses of Daily Symptoms in Women with Fibromyalgia Syndrome. *The Journal Of Pain*, 12(1), 84-93.
- Outcalt, S., Kroenke, K., Krebs, E., Chumbler, N., Wu, J., Yu, Z., & Bair, M. (2015). Chronic pain and comorbid mental health conditions: independent associations of posttraumatic stress disorder and depression with pain, disability, and quality of life. *J Behav Med*, 38(3), 535-543.
- Pagano, T., Matsutani, L., Ferreira, E., Marques, A., & Pereira, C. (2004). Assessment of anxiety and quality of life in fibromyalgia patients. *Sao Paulo Med. J.*, 122(6). <http://dx.doi.org/10.1590/s1516-31802004000600005>

- Perneger, T. (1998). What's wrong with Bonferroni adjustments? *BMJ*, 316(7139), 1236-1238.
- Peterlin, B., Tietjen, G., Brandes, J., Rubin, S., Drexler, E., Lidicker, J., & Meng, S. (2009). Posttraumatic Stress Disorder in Migraine. *Headache: The Journal Of Head And Face Pain*, 49(4), 541-551.
- Pinquart, M. (2012). Self-esteem of children and adolescents with chronic illness: a meta-analysis. *Child: Care, Health And Development*, 39(2), 153-161.
- Pinto-Gouveia, J., Duarte, C., Matos, M., & Fráguas, S. (2013). The Protective Role of Self-compassion in Relation to Psychopathology Symptoms and Quality of Life in Chronic and in Cancer Patients. *Clinical Psychology & Psychotherapy*, 21(4), 311-323.
- Platten, M., Newman, E., & Quayle, E. (2012). Self-Esteem and Its Relationship to Mental Health and Quality of Life in Adults with Cystic Fibrosis. *J Clin Psychol Med Settings*, 20(3), 392-399.
- Preacher, K. & Hayes, A. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*, 40(3), 879-891.
- Purdie, F. & Morley, S. (2015). Self-compassion, pain, and breaking a social contract. *PAIN*, 156(11), 2354-2363.
- Queiroz, L. (2013). Worldwide Epidemiology of Fibromyalgia. *Curr Pain Headache Rep*, 17(8). <http://dx.doi.org/10.1007/s11916-013-0356-5>
- Raes, F., Pommier, E., Neff, K., & Van Gucht, D. (2011). Construction and factorial validation of a short form of the Self-Compassion Scale. *Clinical Psychology & Psychotherapy*, 18(3), 250-255.
- Rosenberg, M. (1965). *Society and the adolescent self-image*. Princeton, N.J.: Princeton University Press.
- Roshan, R., Shariatpanahi, S., Tavoli, A., Qafori, Z., & Fathi-Ashtiani, A. (2008). Quality of life in women with fibromyalgia. *Archives Of Medical Science*, 5, 105-117.
- Rustøen, T., Wahl, A., Hanestad, B., Lerdal, A., Paul, S., & Miaskowski, C. (2004). Gender differences in chronic pain—findings from a population-based study of Norwegian adults. *Pain Management Nursing*, 5(3), 105-117.
- Salaffi, F., Sarzi-Puttini, P., Girolimetti, R., Gasparini, S., Atzeni, F., & Grassi, W. (2009). Development and validation of the self-administered Fibromyalgia Assessment Status: a

- disease-specific composite measure for evaluating treatment effect. *Arthritis Res Ther*, 11(4), R125.
- Sanderson, S., Tatt, I., & Higgins, J. (2007). Tools for assessing quality and susceptibility to bias in observational studies in epidemiology: a systematic review and annotated bibliography. *International Journal Of Epidemiology*, 36(3), 666-676.
- Schaefer, K. (1997). Health patterns of women with fibromyalgia. *J Adv Nurs*, 26(3), 565-571. <http://dx.doi.org/10.1046/j.1365-2648.1997.t01-17-00999.x>
- Scottish Intercollegiate Guidelines Network (SIGN). (2008). *Sign.ac.uk*. Retrieved 1 May 2016, from <http://www.sign.ac.uk/guidelines/fulltext/50/section1.html>
- Segura-Jiménez, V., Álvarez-Gallardo, I., Carbonell-Baeza, A., Aparicio, V., Ortega, F., Casimiro, A., & Delgado-Fernández, M. (2015). Fibromyalgia has a larger impact on physical health than on psychological health, yet both are markedly affected: The al-Ándalus project. *Seminars In Arthritis And Rheumatism*, 44(5), 563-570.
- Shrive, F., Stuart, H., Quan, H., & Ghali, W. (2015). Dealing with missing data in a multi-question depression scale: a comparison of imputation methods. *BMC Medical Research Methodology*, 6, 57.
- Simpson, J., Lekwuwa, G., & Crawford, T. (2013). Illness beliefs and psychological outcome in people with Parkinson's disease. *Chronic Illness*, 9(2), 165-176.
- Sivas, F., Baskan, B., & Aktekin, L. (2009). Depression in fibromyalgia patients' assessment of sleep disorders and quality of life. *Turkish Journal Physical Medical Rehabilitation*, 55, 8-12.
- Skevington, S. & McCrate, F. (2011). Expecting a good quality of life in health: assessing people with diverse diseases and conditions using the WHOQOL-BREF. *Health Expectations*, 15(1), 49-62.
- Smith, B., Papp, Z., Tooley, E., Montague, E., Robinson, A., & Cosper, C. (2010). Traumatic events, perceived stress and health in women with fibromyalgia and healthy controls. *Stress And Health*, 26(1), 83-93
- Sonderen, E., Sanderman, R., & Coyne, J. (2013). Correction: Ineffectiveness of Reverse Wording of Questionnaire Items: Let's Learn from Cows in the Rain. *Plos ONE*, 8(9). <http://dx.doi.org/10.1371/annotation/af78b324-7b44-4f89-b932-e851fe04a8e5>

- Soriano-Maldonado, A., Amris, K., Ortega, F., Segura-Jiménez, V., Estévez-López, F., & Álvarez-Gallardo, I. et al. (2015). Association of different levels of depressive symptoms with symptomatology, overall disease severity, and quality of life in women with fibromyalgia. *Qual Life Res*, 24(12), 2951-2957.
- Stewart, L. A., & Tierney, J. F. (2002). To IPD or not to IPD? Advantages and disadvantages of systematic reviews using individual patient data. *Evaluation & the health professions*, 25, 76-97.
- Tajvar, M., Arab, M., & Montazeri, A. (2008). Determinants of health-related quality of life in elderly in Tehran, Iran. *BMC Public Health*, 8(1), 323.
- Tan, G., Jensen, M., Thornby, J., & Shanti, B. (2004). Validation of the brief pain inventory for chronic nonmalignant pain. *The Journal Of Pain*, 5(2), 133-137.
- Tander, B., Cengiz, K., Alayli, G., İlhanli, İ., Canbaz, S., & Canturk, F. (2008). A comparative evaluation of health related quality of life and depression in patients with fibromyalgia syndrome and rheumatoid arthritis. *Rheumatol Int*, 28(9), 859-865. <http://dx.doi.org/10.1007/s00296-008-0551-6>
- Terry, M. & Leary, M. (2011). Self-compassion, self-regulation, and health. *Self And Identity*, 10(3), 352-362.
- The Scottish Intercollegiate Guidelines Network (SIGN) Management of Chronic Pain. (2013). *SIGN Publication No.136*, 136. Retrieved from <http://sign.ac.uk>
- Timmerman, G., Calfa, N., & Stuifbergen, A. (2013). Correlates of Body Mass Index in Women with Fibromyalgia. *Orthopaedic Nursing*, 32(2), 113-119.
- Torre, F., Martín-Corral, J., Callejo, A., Gómez-Vega, C., La Torre, S., & Esteban, et al. (2008). Calidad de vida relacionada con la salud y estrategias de afrontamiento ante el dolor en pacientes atendidos por una unidad de tratamiento del dolor. *Revista De La Sociedad Española Del Dolor*, 15(2), 83-93.
- Turk, D., Dworkin, R., Trudeau, J., Benson, C., Biondi, D., Katz, N., & Kim, M. (2015). Validation of the Hospital Anxiety and Depression Scale in Patients With Acute Low Back Pain. *The Journal Of Pain*, 16(10), 1012-1021.

- Tüzün, E., Albayrak, G., Eker, L., Sözüay, S., & Daşkapan, A. (2004). A comparison study of quality of life in women with fibromyalgia and myofascial pain syndrome. *Disability And Rehabilitation*, 26(4), 198-202.
- University of York, C. (2008). *Systematic Reviews: CRD's guidance for undertaking reviews in healthcare*. York Publishing Services Ltd. University of York. Retrieved 1 March 2016, from http://www.york.ac.uk/media/crd/Systematic_Reviews.pdf
- Unruh, A., Ritchie, J., & Merskey, H. (1999). Does Gender Affect Appraisal of Pain and Pain Coping Strategies? *The Clinical Journal Of Pain*, 15(1), 31-40.
- van den Beuken-van Everdingen, M., de Rijke, J., Kessels, A., Schouten, H., van Kleef, M., & Patijn, J. (2007). Prevalence of pain in patients with cancer: a systematic review of the past 40 years. *Annals Of Oncology*, 18(9), 1437-1449
- Van Son, M., De Vries, J., Roukema, J., & Den Oudsten, B. (2013). Health status, health-related quality of life, and quality of life following ankle fractures: A systematic review. *Injury*, 44(11), 1391-1402.
- Vandenbroucke, J. (2007). Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and Elaboration. *Annals Of Internal Medicine*, 147(8), W.
- Wallace, B., Dahabreh, I., Trikalinos, T., Lau, J., Trow, P., & Schmid, C. (2012). Closing the Gap between Methodologists and End-Users: R as a Computational Back-End. *Journal Of Statistical Software*, 49(5).
- Ware, J. & Sherbourne, C. (1992). The MOS 36-Item Short-Form Health Survey (SF-36). *Medical Care*, 30(6), 473-483.
- Ware, J., Kosinski, M., & Dewey, J. (2000). *How to score version 2 of the SF-36 health survey (standard & acute forms)*. Lincoln, R.I.: QualityMetric Inc.
- Wetherell, J., Afari, N., Rutledge, T., Sorrell, J., Stoddard, J., & Petkus, A. et al. (2011). A randomized, controlled trial of acceptance and commitment therapy and cognitive-behavioral therapy for chronic pain. *Pain*, 152(9), 2098-2107.
- White, K., Speechley, M., Harth, M., & Ostbye, T. (1999). Comparing self-reported function and work disability in 100 community cases of fibromyalgia syndrome versus controls in London, Ontario: the London Fibromyalgia Epidemiology Study. *Arthritis & Rheumatism*, 42(1), 76-83. [http://dx.doi.org/10.1002/1529-0131\(199901\)42:1<76::aid-anr10>3.0.co;2-g](http://dx.doi.org/10.1002/1529-0131(199901)42:1<76::aid-anr10>3.0.co;2-g)

- Williams, A., Eccleston, C., & Morley, S. (2009). 1012 Systematic Review and meta-analysis of psychological treatments for persistent pain in adults, excluding headache. *European Journal Of Pain*, 13, S284. [http://dx.doi.org/10.1016/s1090-3801\(09\)61015-6](http://dx.doi.org/10.1016/s1090-3801(09)61015-6)
- Williams, M., Dalgleish, T., Karl, A., & Kuyken, W. (2014). Examining the factor structures of the Five Facet Mindfulness Questionnaire and the Self-Compassion Scale. *Psychological Assessment*, 26(2), 407-418.
- Wolfe, F., Clauw, D., Fitzcharles, M., Goldenberg, D., Katz, R., & Mease, P. et al. (2010). The American College of Rheumatology Preliminary Diagnostic Criteria for Fibromyalgia and Measurement of Symptom Severity. *Arthritis Care Res*, 62(5), 600-610.
- Wolfe, F., Michaud, K., Li, T., & Katz, R. (2009). EQ-5D and SF-36 Quality of Life Measures in Systemic Lupus Erythematosus: Comparisons with Rheumatoid Arthritis, Noninflammatory Rheumatic Disorders, and Fibromyalgia. *The Journal Of Rheumatology*, 37(2), 296-304.
- Wolfe, F., Smythe, H., Yunus, M., Benett, R., & Bombardier, C. (2016). The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the Multicentre Criteria Committee. *Arthritis, Rheumatology*, 33, 160-172.
- Wong, W. & Fielding, R. (2011). Prevalence and Characteristics of Chronic Pain in the General Population of Hong Kong. *The Journal Of Pain*, 12(2), 236-245.
- Woolf, C. (2011). Central sensitization: Implications for the diagnosis and treatment of pain. *Pain*, 152(Supplement), S2-S15. <http://dx.doi.org/10.1016/j.pain.2010.09.030>
- Wren, A., Somers, T., Wright, M., Goetz, M., Leary, M., & Fras, A. et al. (2012). Self-Compassion in Patients With Persistent Musculoskeletal Pain: Relationship of Self-Compassion to Adjustment to Persistent Pain. *Journal Of Pain And Symptom Management*, 43(4), 759-770.
- Yarnell, L., Stafford, R., Neff, K., Reilly, E., Knox, M., & Mullarkey, M. (2015). Meta-Analysis of Gender Differences in Self-Compassion. *Self And Identity*, 14(5), 499-520.
- Yu, L., Norton, S., Harrison, A., & McCracken, L. (2015). In search of the person in pain: A systematic review of conceptualization, assessment methods, and evidence for self and identity in chronic pain. *Journal Of Contextual Behavioral Science*, 4(4), 246-262. <http://dx.doi.org/10.1016/j.jcbs.2015.10.003>
- Yunus, M. (2002). Gender differences in fibromyalgia and other related syndromes. *The Journal Of Gender-Specific Medicine*, 5(2), 42-47.

Zigmond, A. & Snaith, R. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, 67, 361-370.



JOURNAL OF CONTEXTUAL BEHAVIORAL SCIENCE

AUTHOR INFORMATION PACK

TABLE OF CONTENTS

• Description	p.1
• Editorial Board	p.2
• Guide for Authors	p.3



ISSN: 2212-1447

DESCRIPTION

The *Journal of Contextual Behavioral Science* is the official journal of the [Association for Contextual Behavioral Science \(ACBS\)](#).

Contextual Behavioral Science is a systematic and pragmatic approach to the understanding of **behavior**, the solution of human problems, and the promotion of human growth and development. Contextual Behavioral Science uses functional principles and theories to analyze and modify action embedded in its historical and situational context. The goal is to predict and influence behavior, with precision, scope, and depth, across all behavioral domains and all levels of analysis, so as to help create a **behavioral science** that is more adequate to the challenge of the human condition.

JCBS welcomes **contextual behavioral analyses** of phenomena that are relevant to the aims and scope of the society's mission, which is to change behavior at an individual or cultural level, to alleviate human suffering, and to advance human wellbeing. JCBS is also a strategic approach to the analysis of human behavior that proposes the need for a multi-level (e.g. social factors, neurological factors, behavioral factors) and multi-method (e.g., time series analyses, cross-sectional, experimental...) exploration of contextual and manipulable variables relevant to the prediction and influence of **human behavior**. In addition it places a strong emphasis in theory development and the promotion of effective practices that link back to scientific principles.

The journal considers papers relevant to a contextual behavioral approach include empirical studies (without topical restriction - e.g., clinical psychology, psychopathology, education, organizational psychology, etc.), reviews (systematic reviews and meta-analyses are preferred), and conceptual and philosophical papers on contextual behavioral science. We are particularly interested in papers emphasizing the study of core behavioral processes that are relevant to a broad range of human problems, and thus not limited to certain populations. Conceptual papers selected for publication may address a broad range of topics but generally will focus on contextual and functional variables or the philosophical analysis of contextual behavioral science. Papers that challenge a contextual behavioral science approach are always welcome. Papers bridging different approaches (e.g., connecting behavioral approaches with cognitive views; or neurocognitive psychology; or evolutionary science) are particularly encouraged.

The journal publishes papers written by researchers, practitioners, and theoreticians from different intellectual traditions. What is distinctive is not a narrowly defined theory or set of applied methods but whether the methodology, conceptualization, or strategy employed is relevant to a contextual behavioral approach.

GUIDE FOR AUTHORS

Types of article

All manuscripts must clearly and explicitly be of relevance to CBS. You may find the JCBS article "[Contextual Behavioral Science: creating a science more adequate to the challenge of the human condition](#)" helpful in assessing whether your manuscript is likely to be of interest to readers of this journal.

Articles should fall into one of seven categories:

1. Empirical research (up to 6000 words)
2. Brief empirical reports (up to 3000 words)
3. Review articles (up to 10,000 words)
4. Conceptual articles (up to 6000 words)
5. In practice (up to 3000 words)
6. Practical innovations (up to 3000 words)
7. Professional interest briefs (up to 3000 words)

Word limits exclude references, tables and figures but include the abstract

1. Empirical research. JCBS welcomes manuscripts across a breadth of domains from basic behavioral science to clinical trials. Research concerning the measurement and testing of process of change is particularly welcome. Potential methodologies include but are not limited to: randomized controlled trials, single case experimental designs, cross-sectional and prospective cohort studies, mixed-methods designs, small scale analog studies. Papers reporting null findings are also welcome if their methodology is sound and their power sufficient. Authors of such papers will need to emphasize the implications of their findings for future research and practice.

2. Brief empirical reports. Manuscripts in this section may report preliminary, provocative or replicated results. Empirically sound methodology and adequate power remain important considerations.

3. Review articles. Manuscripts reviewing a wide range of topics are encouraged as long as their content is directly relevant to CBS. Systematic reviews and meta-analyses are particularly welcome. Authors are advised to consult relevant MARS (www.apa.org/pubs/authors/jars.pdf) and PRISMA resources (<http://www.prisma-statement.org/>) when preparing such manuscripts.

4. Conceptual articles. Manuscripts in this section should address conceptual or theoretical issues relevant to CBS. This may include papers that discuss relevant philosophical assumptions and traditions, or conceptual papers which explore aspects of or inconsistencies in contextual behavioral theory and science.

5. In practice. Manuscripts in this section are designed to make CBS useful to practitioners from a wide variety of areas. Manuscripts must be written in an accessible style and should be easily understood by practitioners who are not experts in research or basic behavioral science. Manuscripts should provide both clear insights for new practitioners as well as stating the questions that remain to be answered by future research.

6. Practical innovations. Manuscripts in this section seek to apply the findings and applications of CBS to under-studied, under-served or novel areas. The scope of these manuscripts is limited only by the journal's broad mission: creating a science more adequate to the challenge of the human condition.

7. Professional interest briefs. Manuscripts in this section highlight professional issues of relevance to those working in the field of CBS. Examples include manuscripts related to training and supervision, assessment methods in professional settings or opinions on contemporary issues.

Contact details for submission

To contact the Editor-in-Chief prior to your submission with any questions, please email ciarochij@gmail.com

BEFORE YOU BEGIN

Ethics in publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see <https://www.elsevier.com/publishingethics> and <https://www.elsevier.com/journal-authors/ethics>.

Human and animal rights

If the work involves the use of human subjects, the author should ensure that the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans, <http://www.wma.net/en/30publications/10policies/b3/index.html>; Uniform Requirements for manuscripts submitted to Biomedical journals, <http://www.icmje.org>. Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

All animal experiments should be carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, EU Directive 2010/63/EU for animal experiments, or the National Institutes of Health guide for the care and use of Laboratory animals (NIH Publications No. 8023, revised 1978) and the authors should clearly indicate in the manuscript that such guidelines have been followed. **All animal studies need to ensure they comply with the ARRIVE guidelines.** More information can be found at <http://www.nc3rs.org.uk/page.asp?id=1357>.

Conflict of interest

All authors are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within three years of beginning the submitted work that could inappropriately influence, or be perceived to influence, their work. See also <https://www.elsevier.com/conflictsofinterest>. Further information and an example of a Conflict of Interest form can be found at: http://service.elsevier.com/app/answers/detail/a_id/286/supporthub/publishing.

Submission declaration and verification

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <https://www.elsevier.com/sharingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. To verify originality, your article may be checked by the originality detection service CrossCheck <https://www.elsevier.com/editors/plagdetect>.

Authorship

All authors should have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be submitted.

Changes to authorship

Authors are expected to consider carefully the list and order of authors **before** submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the **corresponding author**: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed.

Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors **after** the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

Article transfer service

This journal is part of our Article Transfer Service. This means that if the Editor feels your article is more suitable in one of our other participating journals, then you may be asked to consider transferring the article to one of those. If you agree, your article will be transferred automatically on your behalf with no need to reformat. Please note that your article will be reviewed again by the new journal. More information about this can be found here: <https://www.elsevier.com/authors/article-transfer-service>.

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright, see <https://www.elsevier.com/copyright>). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <https://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: please consult <https://www.elsevier.com/permissions>.

For open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' (for more information see <https://www.elsevier.com/OAauthoragreement>). Permitted third party reuse of open access articles is determined by the author's choice of user license (see <https://www.elsevier.com/openaccesslicenses>).

Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. For more information see <https://www.elsevier.com/copyright>.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

Funding body agreements and policies

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some authors may also be reimbursed for associated publication fees. To learn more about existing agreements please visit <https://www.elsevier.com/fundingbodies>.

Open access

This journal offers authors a choice in publishing their research:

Open access

- Articles are freely available to both subscribers and the wider public with permitted reuse.
- An open access publication fee is payable by authors or on their behalf (e.g. by their research funder or institution).

Subscription

- Articles are made available to subscribers as well as developing countries and patient groups through our universal access programs (<https://www.elsevier.com/access>).
- No open access publication fee payable by authors.

Regardless of how you choose to publish your article, the journal will apply the same peer review criteria and acceptance standards.

For open access articles, permitted third party (re)use is defined by the following Creative Commons user licenses:

Creative Commons Attribution (CC BY)

Lets others distribute and copy the article, create extracts, abstracts, and other revised versions, adaptations or derivative works of or from an article (such as a translation), include in a collective work (such as an anthology), text or data mine the article, even for commercial purposes, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, and do not modify the article in such a way as to damage the author's honor or reputation.

Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

The open access publication fee for this journal is **USD 1800**, excluding taxes. Learn more about Elsevier's pricing policy: <http://www.elsevier.com/openaccesspricing>.

Green open access

Authors can share their research in a variety of different ways and Elsevier has a number of green open access options available. We recommend authors see our green open access page for further information (<http://elsevier.com/greenopenaccess>). Authors can also self-archive their manuscripts immediately and enable public access from their institution's repository after an embargo period. This is the version that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and in editor-author communications. Embargo period: For subscription articles, an appropriate amount of time is needed for journals to deliver value to subscribing customers before an article becomes freely available to the public. This is the embargo period and it begins from the date the article is formally published online in its final and fully citable form.

This journal has an embargo period of 24 months.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's WebShop (<http://webshop.elsevier.com/languageediting/>) or visit our customer support site (<http://support.elsevier.com>) for more information.

Informed consent and patient details

Studies on patients or volunteers require ethics committee approval and informed consent, which should be documented in the paper. Appropriate consents, permissions and releases must be obtained where an author wishes to include case details or other personal information or images of patients and any other individuals in an Elsevier publication. Written consents must be retained by the author and copies of the consents or evidence that such consents have been obtained must be provided to Elsevier on request. For more information, please review the *Elsevier Policy on the Use of Images or Personal Information of Patients or other Individuals*, <https://www.elsevier.com/patient-consent-policy>. Unless you have written permission from the patient (or, where applicable, the next of kin), the personal details of any patient included in any part of the article and in any supplementary materials (including all illustrations and videos) must be removed before submission.

Submission

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

Referees

Please submit the names and institutional e-mail addresses of several potential referees. For more details, visit our [Support site](#). Note that the editor retains the sole right to decide whether or not the suggested reviewers are used.

PREPARATION

Use of word processing software

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the Guide to Publishing with Elsevier: <https://www.elsevier.com/guidepublication>). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

Article structure

Subdivision - unnumbered sections

Divide your article into clearly defined sections. Each subsection is given a brief heading. Each heading should appear on its own separate line. Subsections should be used as much as possible when cross-referencing text: refer to the subsection by heading as opposed to simply 'the text'.

Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

Material and methods

Provide sufficient detail to allow the work to be reproduced. Methods already published should be indicated by a reference: only relevant modifications should be described.

Theory/calculation

A Theory section should extend, not repeat, the background to the article already dealt with in the Introduction and lay the foundation for further work. In contrast, a Calculation section represents a practical development from a theoretical basis.

Results

Results should be clear and concise.

Discussion

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

Appendices

If there is more than one appendix, they should be identified as A, B, etc. Formulae and equations in appendices should be given separate numbering: Eq. (A.1), Eq. (A.2), etc.; in a subsequent appendix, Eq. (B.1) and so on. Similarly for tables and figures: Table A.1; Fig. A.1, etc.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.

- **Author names and affiliations.** Please clearly indicate the given name(s) and family name(s) of each author and check that all names are accurately spelled. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.

- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.**

• **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Graphical abstract

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. See <https://www.elsevier.com/graphicalabstracts> for examples.

Authors can make use of Elsevier's Illustration and Enhancement service to ensure the best presentation of their images and in accordance with all technical requirements: [Illustration Service](#).

Highlights

Highlights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). See <https://www.elsevier.com/highlights> for examples.

Keywords

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Formatting of funding sources

List funding sources in this standard way to facilitate compliance to funder's requirements:

Funding: This work was supported by the National Institutes of Health [grant numbers xxxx, yyyy]; the Bill & Melinda Gates Foundation, Seattle, WA [grant number zzzz]; and the United States Institutes of Peace [grant number aaaa].

It is not necessary to include detailed descriptions on the program or type of grants and awards. When funding is from a block grant or other resources available to a university, college, or other research institution, submit the name of the institute or organization that provided the funding.

If no funding has been provided for the research, please include the following sentence:

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Math formulae

Please submit math equations as editable text and not as images. Present simple formulae in line with normal text where possible and use the solidus (/) instead of a horizontal line for small fractional terms, e.g., X/Y . In principle, variables are to be presented in italics. Powers of e are often more conveniently denoted by \exp . Number consecutively any equations that have to be displayed separately from the text (if referred to explicitly in the text).

Footnotes

Footnotes should be used sparingly. Number them consecutively throughout the article. Many word processors can build footnotes into the text, and this feature may be used. Otherwise, please indicate the position of footnotes in the text and list the footnotes themselves separately at the end of the article. Do not include footnotes in the Reference list.

Artwork

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the published version.
- Submit each illustration as a separate file.

A detailed guide on electronic artwork is available on our website:

<https://www.elsevier.com/artworkinstructions>.

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. For further information on the preparation of electronic artwork, please see <https://www.elsevier.com/artworkinstructions>.

Figure captions

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

References in a special issue

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

Reference management software

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support Citation Style Language styles (<http://citationstyles.org>), such as Mendeley (<http://www.mendeley.com/features/reference-manager>) and Zotero (<https://www.zotero.org/>), as well as EndNote (<http://endnote.com/downloads/styles>). Using the word processor plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide.

Users of Mendeley Desktop can easily install the reference style for this journal by clicking the following link:

<http://open.mendeley.com/use-citation-style/journal-of-contextual-behavioral-science>

When preparing your manuscript, you will then be able to select this style using the Mendeley plug-ins for Microsoft Word or LibreOffice.

Reference style

Text: Citations in the text should follow the referencing style used by the American Psychological Association. You are referred to the Publication Manual of the American Psychological Association, Sixth Edition, ISBN 978-1-4338-0561-5, copies of which may be ordered from <http://books.apa.org/books.cfm?id=4200067> or APA Order Dept., P.O.B. 2710, Hyattsville, MD 20784, USA or APA, 3 Henrietta Street, London, WC3E 8LU, UK.

List: references should be arranged first alphabetically and then further sorted chronologically if necessary. More than one reference from the same author(s) in the same year must be identified by the letters 'a', 'b', 'c', etc., placed after the year of publication.

Examples:

Reference to a journal publication:

Van der Geer, J., Hanraads, J. A. J., & Lupton, R. A. (2010). The art of writing a scientific article. *Journal of Scientific Communications*, 163, 51–59.

Reference to a book:

Strunk, W., Jr., & White, E. B. (2000). *The elements of style*. (4th ed.). New York: Longman, (Chapter 4).

Reference to a chapter in an edited book:

Mettam, G. R., & Adams, L. B. (2009). How to prepare an electronic version of your article. In B. S. Jones, & R. Z. Smith (Eds.), *Introduction to the electronic age* (pp. 281–304). New York: E-Publishing Inc.

Reference to a website:

Cancer Research UK. Cancer statistics reports for the UK. (2003). <http://www.cancerresearchuk.org/aboutcancer/statistics/cancerstatsreport/> Accessed 13.03.03.

Video data

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the files in one of our recommended file formats with a preferred maximum size of 150 MB. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our video instruction pages at <https://www.elsevier.com/artworkinstructions>. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

Supplementary material

Supplementary material can support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high-resolution images, background datasets, sound clips and more. Please note that such items are published online exactly as they are submitted; there is no typesetting involved (supplementary data supplied as an Excel file or as a PowerPoint slide will appear as such online). Please submit the material together with the article and supply a concise and descriptive caption for each file. If you wish to make any changes to supplementary data during any stage of the process, then please make sure to provide an updated file, and do not annotate any corrections on a previous version. Please also make sure to switch off the 'Track Changes' option in any Microsoft Office files as these will appear in the published supplementary file(s). For more detailed instructions please visit our artwork instruction pages at <https://www.elsevier.com/artworkinstructions>.

AudioSlides

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. More information and examples are available at <https://www.elsevier.com/audioslides>. Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

Submission checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded, and contain:

- Keywords
- All figure captions
- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
- References are in the correct format for this journal
- All references mentioned in the Reference list are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)

Printed version of figures (if applicable) in color or black-and-white

- Indicate clearly whether or not color or black-and-white in print is required.

For any further information please visit our customer support site at <http://support.elsevier.com>.

AFTER ACCEPTANCE

Online proof correction

Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

Offprints

The corresponding author will, at no cost, receive a customized [Share Link](#) providing 50 days free access to the final published version of the article on [ScienceDirect](#). The Share Link can be used for sharing the article via any communication channel, including email and social media. For an extra charge, paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's [Webshop](#). Corresponding authors who have published their article open access do not receive a Share Link as their final published version of the article is available open access on ScienceDirect and can be shared through the article DOI link.

AUTHOR INQUIRIES

You can track your submitted article at <https://www.elsevier.com/track-submission>. You can track your accepted article at <https://www.elsevier.com/trackarticle>. You are also welcome to contact Customer Support via <http://support.elsevier.com>.

© Copyright 2014 Elsevier | <http://www.elsevier.com>

Appendix B- Quality Rating Tool for Systematic Review

	Yes	Partial	No
Sampling: The recruitment method			
The recruitment/sampling method ensures that minimal bias is introduced (e.g. recruitment from the general population)			
The recruitment/sampling method used random sampling			
The recruitment/sampling used a convenience sample			
Validated Method for ascertaining severity of Fibromyalgia			
The psychometric properties of the outcome measure for fibromyalgia are valid and reliable in this study population			
The measure has not been validated in this population			
The measure has not undergone any psychometric evaluation, or did not used a measure to assess severity of fibromyalgia			
Power and Sample Size			
The study has a sample size large enough to detect a small to moderate effect size with the statistical power of 0.8 at an alpha level of 0.05			
The study has a sample size large enough to detect a small to moderate effect size the statistical power of 0.7 at an alpha level of 0.05			
The study has a sample size large enough to detect a small to moderate effect size with the statistical power of less than 0.7 at an alpha level of 0.05 or power was not calculable			
Appropriate Method used			
The method of statistical analysis was appropriate to the research question being asked. Confidence intervals, p-values and effect sizes are reported.			
The analysis is appropriate however the findings are not reported in sufficient detail			
The method of analysis is not appropriate to the research question and does not provide meaningful results			
Confounding variables			
Potential confounding variables are adequately recognised and a description is provided on how they are addressed in statistical analysis			
Potential confounding variables are recognised and there is recognition of a possible effect but they are not considered in statistical analysis			
Potential confounding variables are not considered			

Appendix C- Excluded Studies from Systematic Review

Studies excluded	Reasons for exclusion
Full Article Review	
Alves, 2012	No control
Amital 2014	Included both FM and Systemic sclerosis in the sample
Andrell 2014	SF—36 were z scores
Angst 2006	Needed to have had history of failed treatment to participate in the study
Aparicio 2013	Did not include SF-36 subscales
Aparicio 2014	Inclusion criteria obesity
Arranz 2012	Split FM according to BMI
Assis, 2006	No control
Atagun, 2012	Duplicate
Berfman, 2005	Did not include SF-36 subscales
Boehm, 2011	Did not include SF-36 subscales
Borsbo, 2010	Did not include SF-36 subscales
Brattberg, 2008	Had to be non-working
Brill, 2012	Did not include SD for SF-36 means
Burckhardt, 2005	Did not include SF-36 subscales
Buskila, 2000	Did not include SF-36 subscales
Campos, 2012	No control
Campos, 2013	Split FM into age cohorts
Carbonell-Baeza, 2011	Did not include SF-36 subscales
Cardona-Arias, 2014	No Controls
Castelli, 2012	No Controls
Chen, 2014	Conference abstract- unavailable
Choy, 2009	Did not include SF-36 subscales
Coster, 2008	Control group included people with pain
DaCosta, 2000	Could not access article
Danis, 2015	Abstract conference- unavailable
De Araujo	No control group
Dell’Osso, 2011	Could not access article
Dogramaci, 2009	Split FM into skin conditions
Donaldson, 2001	No Control
Glattacker, 2010	Treatment Groups
Jensen, 2009	Did not include SF-36 subscales
Jiao, 2015	Sample from a previous treatment group
Jiao, 2015	Split FM into history of abuse
Jiao, 2014	Divided into age cohorts
Kim, 2012	Split FM into BMI
Kim, 2012	FM in BMI
Kim, 2013	Same data set
Kim, 2012	Did not include SF-36 subscales
Latorre, 2013	Treatment study
Martinez, 2001	No standard deviation for the SF-36 subscales
Matsumoto, 2011	No control
Oliveiria, 2009	No control
Oncu, 2013	RA control group

Peleg, 2008	No control
Picavet, 2003	No control
Raj, 2000	No control
Rodero, 2010	No control
Romero, 2012	Treatment control
Rosenzweig, 2010	No control
Sanudo, 2011	RCT- treatment
Sanudo, 2012	RCT- same data as above
Saxe, 2009	Did not include SF-36 subscales
Schlenck, 1998	Did not include SF-36 subscales
Schlenck, 1998	Used a CBT treatment groups
Seo, 2010	Abstract Conference- unavailable
Servant, 2011	Did not include SF-36 subscales
Sigl-Erkel, 2011	RCT
Silvas, 2009	RA control
Slawsky, 2011	Used the Rand SF-36
Taggart,	No control
Theadorn, 2007	No control
Trinanes, 2015	No control
Tomas-Carus, 2009	No control
Trovato, 2010	Did not include control
Turan, 2011	No control
Uhlemann, 2007	Did not include control
Ulas, 2006	Duplicate datas
Valim, 2002	Unable to access article
Verbunt, 2008	Did not include SF-36 subdomains
Vitorino, 2006	No Control
Yilmaz, 2008	No Control
Yoshikawa, 2010	Had to meet clinical cut-off for depression
Walker, 1997	No Control
Wang, 2010	RCT-did not include subdomains
Wang, 2010	Did not include SF-36 subdomains
Wolfe, 2013	Abstract Conference- Unavailable
Wolfe, 2011	No control
Wolfe, 2005	No control
Wolfe, 2009	No control
Wolfe, 2010	No control
Wolfe, 2011	No control
Wong, 2012	RCT-did not include subdomains
Web of Science	
Efrati, 2015	Did not include SF-36 subdomains
Plazier, 2015	Did not include SF-36 subdomains
Wolfe, 2014	Did not include SF-36 subdomains
Pedro, 2012	Split FM into severity
Rombaut, 2011	Did not include SF-36 subdomains
Ubago, 2007	Did not include SF-36
Mease, 2008	Did not include SF-36 subdomains
Kaziyama, 2005	Abstract Conference- unavailable

Appendix D: Systematic Review Data Extraction Form

General Information

Date of data extraction

Author

Article Title

Citation

Type of publication (e.g. journal articles, conference abstract)

Country of origin

Study Characteristics

Aim/objectives of the study

Study design

Study inclusion criteria

Study exclusion criteria

Recruitment procedures used

Participant Characteristics

Age

Gender

Ethnicity

Socio-economic status

Mean \pm SD SF-36 subscales

Mean Fibromyalgia severity

Mean depression

Co-morbidities

Disease Duration

Number of participants enrolled in study

Number of participants who provided valid data

Method

Scoring and use of SF-36

FM measure

Depression measure

Setting in which the study is conducted

Description of the controls

Outcome data/results

Additional measures/confounding variables


Statistical analysis used

Results of study analysis


Key Findings

Appendix E- Forest Plots Meta-Analysed Pooled Mean Scores for FM for SF-36 subdomains


Physical Function




Physical Role




Bodily Pain




General Health




Vitality




Social Functioning



Emotion Role




Mental Health




Appendix F-Difference Between Means for FM and Control Group on SF-36 Subdomains


Physical Function




Physical Role




Bodily Pain




General Health




Vitality




Social Functioning



Emotional Role



Mental Health



Appendix G - Sensitivity and Subgroup Analysis Table

Sensitivity Analysis	PF	PR	BP	GH	VT	SF	ER	MH
Primary Analysis	-33.9 (-40.3,-29.5) I ² = 95.9 18 studies	-53.4 (-62.9,-43.9) I ² = 97.9 18 studies	-38.2(-44.0,-32.5) I ² = 96.9 18 studies	-31.1(-35.9,-26.3) I ² = 95.5 18 studies	-29.4(-34.6,-24.2) I ² = 96.4 18 studies	-24.4(-30.5,-18.2) I ² = 96.8 18 studies	-37.5(-45.9,-29.0) I ² = 96.3 18 studies	-21.2(-24.8,-17.7) I ² = 91.6 18 studies
Excluding convenience, non-randomised or unreported sampling methods	-38.6(-43.7,-33.6) I ² = 89.5 4 Studies	-56.8(-72.1,-41.4) I ² = 98.4 4 Studies	-45.3(-49.2,-41.5) I ² = 11.1 4 Studies	-35.2(-44.6,-25.8) I ² = 97.3 4 Studies	-38.2(-45.0,-31.4) I ² = 94.9 4 Studies	-35.4(-50.7,-20.1) I ² = 98.6 4 Studies	-36.6(-45.4,-27.8) I ² = 90.0 4 Studies	-23.7(-25.1,-22.3) I ² = 0 4 Studies
Assessed Severity of FM >39	-35.8(-43.3,-28.3) I ² = 92.2 6 Studies	-50.6(-60.2,-41.0) I ² = 93.6 6 Studies	-39.2(-48.9,-29.5) I ² = 96.2 6 Studies	-29.9(-37.5,-22.2) I ² = 92.3 6 Studies	-30.9(-41.6,-20.3) I ² = 96.9 6 Studies	-20.7(-31.4,-10.00) I ² = 96.0 6 Studies	-38.1(-53.2,-23.0) I ² = 96.5 6 Studies	-22.3(-29.6,-15.0) I ² = 93.0 6 Studies
Excluding studies considered to be under power	-33.8(-43.0,-24.6) I ² = 97.9 8 Studies	-46.7(-61.2,-32.2) I ² = 98.7 8 Studies	-39.4(-49.0,-29.8) I ² = 98.4 8 Studies	-29.3(-37.6,-21.1) I ² = 97.6 8 Studies	-32.0(-40.1,-23.8) I ² = 97.8 8 Studies	-22.4(-33.0,-11.9) I ² = 98.3 8 Studies	-31.0(-42.4,-19.4) I ² = 97.2 8 Studies	-20.7(-26.1,-15.3) I ² = 94.3 8 Studies
Excluding studies that did not report findings in sufficient detail	-39.2(-45.8,-32.6) I ² = 90.3 4 Studies	-45.2(-57.2,-33.2) I ² = 95.6 4 Studies	-47.6(-55.4,-39.7) I ² = 93.7 4 Studies	-34.5(-43.4,-25.7) I ² = 92.9 4 Studies	-34.7(-41.8,-27.5) I ² = 90.6 4 Studies	-16.8(-28.9,-4.72) I ² = 96.7 4 Studies	-36.8(-57.0,-16.6) I ² = 97.7 4 Studies	-25.7(-40.0,-18.5) I ² = 90.5 4 Studies
Excluded studies that did not consider confounding variables in the analysis	-35.7(-40.9,-30.4) I ² = 93.3 12 Studies	-54.8(-64.4,-45.2) I ² = 97.1 12 Studies	-37.1(-44.6,-29.6) I ² = 97.6 12 Studies	-31.9(-36.5,-27.4) I ² = 92.7 12 Studies	-31.7(-37.3,-26.1) I ² = 95.7 12 Studies	-24.1(-30.6,-17.5) I ² = 96.3 12 Studies	-43.0(-52.6,-33.3) I ² = 96.0 12 Studies	-23.5(-28.2,-18.9) I ² = 93.3 12 Studies

Area of Origin Turkey (other)	-33.0 (-40.1;- 25.8) I ² = 87.5 8 Studies	-57.6(-67.0;- 48.2) I ² = 86.7 8 Studies	-37.2(- 47.5;-26.9) I ² = 94.9 8 Studies	-32.4(- 39.9;-25.0) I ² = 90.5 8 Studies	-29.2(-38.2;- 20.2) I ² = 93.2 8 Studies	-18.4(-26.9;- 9.9) I ² = 91.6 8 Studies	-50.0 (-66.1;- 33.8) I ² = 96.3 8 Studies	-24.6(-33.3;- 15.8) I ² = 93.8 8 Studies
Iran	-9.9(-13.0;- 6.8) I ² = 0 2 Studies	-14.80(- 18.7;-11.1) I ² = 0 2 Studies	-7.8(-40.9;- 25.3) I ² = 98.7 2 Studies	-9.2(-14.1;- 4.3) I ² = 50.1 2 Studies	-13.5(-15.9;- 11.0) I ² = 0 2 Studies	-0.2(-8.8;- 8.4) I ² = 80.2 2 Studies	-13.2(-16.2;- 10.2) I ² = 0 2 Studies	-8.5(-15.0;- 2.0) I ² = 73.7 2 Studies
America/South America	-47.6(-54.4;- 40.8) I ² = 49.4 4 Studies	-68.8(-82.4;- 55.1) I ² = 76.3 4 Studies	-46.3(- 52.6;-40.1) I ² = 43.3 4 Studies	-37.3(- 49.4;-25.3) I ² = 83.7 4 Studies	-37.3(-49.4;- 25.3) I ² = 83.7 4 Studies	-33.9(-52.8;- 14.9) I ² = 90.4 4 Studies	-40.5(-61.3;- 19.8) I ² = 79.0 4 Studies	-24.2(-31.3;- 17.1) I ² = 49.3 4 Studies
Central Europe	-38.8(-43.5;- 34.0) I ² = 88.4 4 Studies	-49.4(-64.8;- 34.1) I ² = 98.3 4 Studies	-45.3(- 49.2;-41.3) I ² = 86.3 4 Studies	-33.8(- 42.6;-25.0) I ² = 97.0 4 Studies	-38.8(-45.6;- 31.9) I ² = 95.3 4 Studies	-32.1(-45.8;- 18.5) I ² = 98.2 4 Studies	-29.3(-35.4;- 23.3) I ² = 82.7 4 Studies	-23.6(-25.0;- 22.2) I ² = 0 4 Studies

A

The Clinical Journal of Pain



Author Resources

[Instructions for Authors \(this page\)](#)

[Copyright Transfer \(PDF\)](#)

[Reprint Ordering](#)

[Permissions Requests](#)

[Reprints](#)

INSTRUCTIONS FOR AUTHORS

The Clinical Journal of Pain publishes original articles in the following forms: *Clinical investigations*: Present results of original clinical research. *Case reports*: **Case reports will no longer be accepted for publication in Clinical Journal of Pain and thus no submission for case reports will be accepted as of June 13, 2013.** *Reviews*: Comprehensive surveys covering a broad area. They consolidate old ideas and may suggest new ones. They must provide a critique of the literature. *Special articles*: On subjects not easily classified above (e.g., articles on history, education, demography, ethics, socioeconomics, etc.). *Letters to the editor*: These may offer criticism of published material, but must be objective, constructive, and educational. A few references, a small table, or relevant illustrations may be used.

Ethical/Legal Considerations: A submitted manuscript must be an original contribution not previously published (except as an abstract or preliminary report), must not be under consideration for publication elsewhere, and, if accepted, must not be published elsewhere in similar form, in any language, without the consent of Lippincott Williams & Wilkins. Each person listed as an author is expected to have participated in the study to a significant extent. Although the editors and referees make every effort to ensure the validity of published manuscripts, the final responsibility rests with the authors, not with the Journal, its editors, or the publisher. **All manuscripts must be submitted on-line through the journal's Web site at <http://cjp.edmgr.com>.** See submission instructions under "On-line manuscript submission."

Patient anonymity and informed consent: It is the author's responsibility to ensure that a patient's anonymity be carefully protected and to verify that any experimental investigation with human subjects reported in the manuscript was performed with informed consent and following all the guidelines for experimental investigation with human subjects required by the institution(s) with which all the authors are affiliated. Authors should mask patients' eyes and remove patients' names from

figures unless they obtain written consent from the patients and submit written consent with the manuscript.

Conflicts of interest: Authors must state all possible conflicts of interest in the manuscript, including financial, consultant, institutional and other relationships that might lead to bias or a conflict of interest. If there is no conflict of interest, this should also be explicitly stated as none declared. All sources of funding should be acknowledged in the manuscript. All relevant conflicts of interest and sources of funding should be included on the title page of the manuscript with the heading "Conflicts of Interest and Source of Funding:". For example:

Conflicts of Interest and Source of Funding: A has received honoraria from Company Z. B is currently receiving a grant (#12345) from Organization Y, and is on the speaker's bureau for Organization X – the CME organizers for Company A. For the remaining authors none were declared.

Copyright: In addition, each author must complete and submit the journal's copyright transfer agreement, which includes a section on the disclosure of potential conflicts of interest based on the recommendations of the International Committee of Medical Journal Editors, "Uniform Requirements for Manuscripts Submitted to Biomedical Journals" (www.icmje.org/update.html).

A copy of the form is made available to the submitting author within the Editorial Manager submission process. Co-authors will automatically receive an Email with instructions on completing the form upon submission.

Compliance with NIH and Other Research Funding Agency Accessibility Requirements

A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW will identify to the National Library of Medicine (NLM) articles that require deposit and will transmit the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Wellcome Trust, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The Copyright Transfer Agreement provides the mechanism.

Permissions: Authors must submit written permission from the copyright owner (usually the publisher) to use direct quotations, tables, or illustrations that have appeared in copyrighted form elsewhere, along with complete details about the source. Any permissions fees that might be required by the copyright owner are the responsibility of the authors requesting use of the borrowed material, not the responsibility of Lippincott Williams & Wilkins.

MANUSCRIPT SUBMISSION

On-line manuscript submission: All manuscripts must be submitted on-line through the Web site: <http://cjp.edmgr.com>. **First-time users:** Please click the Register button from the menu and enter the requested information. On successful registration, you will be sent an e-mail indicating your user name and password. Print a copy of this information for future reference. **Note:** If you have received an e-mail from us with an assigned user ID and password, or if you are a repeat user, do not register again. Just log in. Once you have an assigned ID and password, you do not have to re-register, even if your status changes (that is, author, reviewer, or editor). You can change your

username/password at any time by clicking "Update My Information" at the top of any page in Editorial Manager. **Authors:** Please click the log-in button from the menu at the top of the page and log into the system as an Author. Submit your manuscript according to the author instructions. You will be able to track the progress of your manuscript through the system. If you experience any problems, please contact the Managing Editor, Julie Porter: JuliePorter529@gmail.com or 303-870-6334 (phone).

Preparation of Manuscript

Cover Letter. With your manuscript, please submit a brief cover letter describing your manuscript and provide the names and e-mail addresses of 3-4 suggested reviewers. These should be people who are knowledgeable of the topic of the manuscript and who will not have a conflict of interest serving as reviewers. The Editors may or may not enlist these suggested reviewers.

Manuscripts that do not adhere to the following instructions will be returned to the corresponding author for technical revision before undergoing peer review.

General format: Submit manuscripts in English as a Word file. Double space all copy, including legends, footnotes, tables, and references.

Title page: Include on the title page (a) complete manuscript title; (b) authors' full names, highest academic degrees, and affiliations; (c) name and address for correspondence, including fax number, telephone number, and e-mail address; (d) address for reprints if different from that of corresponding author; and (e) sources of support that require acknowledgment.

The title page must also include disclosure of funding received for this work from any of the following organizations: National Institutes of Health (NIH); Wellcome Trust; Howard Hughes Medical Institute (HHMI); and other(s).

Structured abstract and key words: Limit the abstract to 250 words. Do not cite references in the abstract. Limit the use of abbreviations and acronyms. Use the following subheads: Objectives, Methods, Results, and Discussion. List three to five key words.

Text: Organize the manuscript into four main headings: Introduction, Materials and Methods, Results, and Discussion. Define abbreviations at first mention in text and in each table and figure. If a brand name is cited, supply the manufacturer's name and address (city and state/country). Acknowledge all forms of support, including pharmaceutical and industry support, in an Acknowledgments paragraph.

The Clinical Journal of Pain does not have a required number of words for the text. Please treat your subject thoroughly but not excessively. Perusing several back issues to familiarize yourself with typical accepted article length is recommended.

Abbreviations: For a list of standard abbreviations, consult the *Council of Biology Editors Style Guide* (available from the Council of Science Editors, 9650 Rockville Pike, Bethesda, MD 20814) or other standard sources. Write out the full term for each abbreviation at its first use unless it is a standard unit of measure.

References: The authors are responsible for the accuracy of the references. References should be cited by number in order of citation in the text. Key the references (double-spaced) at the end of the manuscript, in numbered order. Cite unpublished data, such as papers submitted but not yet accepted for publication or personal communications, in parentheses in the text (H. E. Marman, M.D., unpublished data, February, 1997). If there are more than three authors, name only the first three authors and then use et al. Refer to the *List of Journals Indexed in Index Medicus* for abbreviations of journal names, or access the list at <http://www.nlm.nih.gov/tsd/serials/lji.html>. Sample references are given below:

Journal article

1. Prager J, Jacobs M, Johnson KJ. Evaluation of patients for implantable pain modalities: medical and behavioral assessment. *Clin J Pain* 2001;17:206-214.

Book chapter

2. Todd VR. Visual information analysis: frame of reference for visual perception. In: Kramer P, Hinojosa J, eds. *Frames of reference for pediatric occupational therapy*. Philadelphia: Lippincott Williams & Wilkins, 1999:205-256.

Entire book

3. Kellman RM, Marentette LJ. *Atlas of craniomaxillofacial fixation*. Philadelphia: Lippincott Williams & Wilkins, 1999.

Software

4. *Epi Info* [computer program]. Version 6. Atlanta: Centers for Disease Control and Prevention; 1994.

Online journals

5. Friedman SA. Preeclampsia: a review of the role of prostaglandins. *Obstet Gynecol* [serial online]. January 1988;71:22-37. Available from: BRS Information Technologies, McLean, VA. Accessed December 15, 1990.

Database

6. CANCERNET-PDQ [database online]. Bethesda, MD: National Cancer Institute; 1996. Updated March 29, 1996.

World Wide Web

7. Gostin LO. Drug use and HIV/AIDS [JAMA HIV/AIDS Web site]. June 1, 1996. Available at: <http://www.ama-assn.org/special/hiv/ethics>. Accessed June 26, 1997.

Figures:

A) Creating Digital Artwork

1. Learn about the publication requirements for Digital Artwork: <http://links.lww.com/ES/A42>
2. Create, Scan and Save your artwork and compare your final figure to the Digital Artwork Guideline Checklist (below).
3. Upload each figure to Editorial Manager in conjunction with your manuscript text and tables.

B) Digital Artwork Guideline Checklist

Here are the basics to have in place before submitting your digital artwork:

- Artwork should be saved as TIFF, EPS, or MS Office (DOC, PPT, XLS) files. High resolution PDF files are also acceptable.
- Crop out any white or black space surrounding the image.
- Diagrams, drawings, graphs, and other line art must be vector or saved at a resolution of at least 1200 dpi. If created in an MS Office program, send the native (DOC, PPT, XLS) file.
- Photographs, radiographs and other halftone images must be saved at a resolution of at least 300 dpi.
- Photographs and radiographs with text must be saved as postscript or at a resolution of at least 600 dpi.
- Each figure must be saved and submitted as a separate file. Figures should not be embedded in the manuscript text file.

Remember:

- Cite figures consecutively in your manuscript.
- Number figures in the figure legend in the order in which they are discussed.
- Upload figures consecutively to the Editorial Manager web site and enter figure numbers consecutively in the Description field when uploading the files.

Figure legends: Legends must be submitted for all figures. They should be brief and specific, and they should appear on a separate manuscript page after the references. Use scale markers in the image for electron micrographs, and indicate the type of stain used.

Color figures: The journal accepts for publication color figures that will enhance an article. Authors who submit color figures will receive an estimate of the cost for color reproduction. If they decide not to pay for color reproduction, they can request that the figures be converted to black and white at no charge.

Tables: Cite tables consecutively in the text, and number them in that order. Key each on a separate sheet, and include the table title, appropriate column heads, and explanatory legends (including definitions of any abbreviations used). Do not embed tables within the body of the manuscript. They should be self-explanatory and should supplement, rather than duplicate, the material in the text.

Supplemental Digital Content

Supplemental Digital Content (SDC): Authors may submit SDC via Editorial Manager to LWW journals that enhance their article's text to be considered for online posting. SDC may include standard media such as text documents, graphs, audio, video, etc. On the Attach Files page of the submission process, please select Supplemental Audio, Video, or Data for your uploaded file as the Submission Item. If an article with SDC is accepted, our production staff will create a URL with the SDC file. The URL will be placed in the call-out within the article. SDC files are not copy-edited by

LWW staff, they will be presented digitally as submitted. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

SDC Call-outs

Supplemental Digital Content must be cited consecutively in the text of the submitted manuscript. Citations should include the type of material submitted (Audio, Figure, Table, etc.), be clearly labeled as "Supplemental Digital Content," include the sequential list number, and provide a description of the supplemental content. All descriptive text should be included in the call-out as it will not appear elsewhere in the article.

Example:

We performed many tests on the degrees of flexibility in the elbow (see Video, Supplemental Digital Content 1, which demonstrates elbow flexibility) and found our results inconclusive.

List of Supplemental Digital Content

A listing of Supplemental Digital Content must be submitted at the end of the manuscript file. Include the SDC number and file type of the Supplemental Digital Content. This text will be removed by our production staff and not be published.

Example:

Supplemental Digital Content 1.wmv

SDC File Requirements

All acceptable file types are permissible up to 10 MBs. For audio or video files greater than 10 MBs, authors should first query the journal office for approval. For a list of all available file types and detailed instructions, please visit <http://links.lww.com/A142>.

Style: Pattern manuscript style after the *American Medical Association Manual of Style* (9th edition). *Stedman's Medical Dictionary* (27th edition) and *Merriam Webster's Collegiate Dictionary* (10th edition) should be used as standard references. Refer to drugs and therapeutic agents by their accepted generic or chemical names, and do not abbreviate them. Use code numbers only when a generic name is not yet available. In that case, supply the chemical name and a figure giving the chemical structure of the drug. Capitalize the trade names of drugs and place them in parentheses after the generic names. To comply with trademark law, include the name and location (city and state in USA; city and country outside USA) of the manufacturer of any drug, supply, or equipment mentioned in the manuscript. Use the metric system to express units of measure and degrees Celsius to express temperatures, and use SI units rather than conventional units.

The editorial office will acknowledge receipt of your manuscript and will give you a manuscript number for reference. Address all inquiries regarding manuscripts not yet accepted or published to the Journal's editorial office: JuliePorter529@gmail.com.

Open access

LWW's hybrid open access option is offered to authors whose articles have been accepted for publication. With this choice, articles are made freely available online immediately upon publication. Authors may take advantage of the open access option at the point of acceptance to ensure that this choice has no influence on the peer review and acceptance process. These articles are subject to the journal's standard peer-review process and will be accepted or rejected based on their own merit.

Authors of accepted peer-reviewed articles have the choice to pay a fee to allow perpetual unrestricted online access to their published article to readers globally, immediately upon publication. The article processing charge for The Clinical Journal of Pain is \$2,700. The article processing charge for authors funded by the Research Councils UK (RCUK) is \$3,375. The publication fee is charged on acceptance of the article and should be paid within 30 days by credit card by the author, funding agency or institution. Payment must be received in full for the article to be published open access.

Authors retain copyright

Authors retain their copyright for all articles they opt to publish open access. Authors grant LWW a license to publish the article and identify itself as the original publisher.

Creative Commons license

Articles opting for open access will be freely available to read, download and share from the time of publication. Articles are published under the terms of the Creative Commons License Attribution-NonCommercial No Derivative 3.0 which allows readers to disseminate and reuse the article, as well as share and reuse of the scientific material. It does not permit commercial exploitation or the creation of derivative works without specific permission. To view a copy of this license visit: <http://creativecommons.org/licenses/by-nc-nd/3.0>.

Compliance with NIH, RCUK, Wellcome Trust and other research funding agency accessibility requirements

A number of research funding agencies now require or request authors to submit the post-print (the article after peer review and acceptance but not the final published article) to a repository that is accessible online by all without charge. As a service to our authors, LWW identifies to the National Library of Medicine (NLM) articles that require deposit and transmits the post-print of an article based on research funded in whole or in part by the National Institutes of Health, Howard Hughes Medical Institute, or other funding agencies to PubMed Central. The revised Copyright Transfer Agreement provides the mechanism. LWW ensures that authors can fully comply with the public access requirements of major funding bodies worldwide. Additionally, all authors who choose the open access option will have their final published article deposited into PubMed Central.

RCUK and Wellcome funded authors can choose to publish their paper as open access with the payment of an article process charge (gold route), or opt for their accepted manuscript to be deposited (green route) into PMC with an embargo.

With both the gold and green open access options, the author will continue to sign the Copyright Transfer Agreement (CTA) as it provides the mechanism for LWW to ensure that the author is fully compliant with the requirements. After signature of the CTA, the author will then sign a License to Publish where they will then own the copyright. Those authors who wish to publish their article via the gold route will be able to publish under the terms of the Attribution 3.0 (CCBY) License. To view a copy of this license visit: <http://creativecommons.org/licenses/by/2.0/>. Those authors who wish to publish their article via the green route will be able to publish under the rights of the Attribution Non-commercial 3.0 (CCBY NC) license (<http://creativecommons.org/licenses/by-nc/2.0/>).

It is the responsibility of the author to inform the Editorial Office and/or LWW that they have RCUK funding. LWW will not be held responsible for retroactive deposits to PMC if the author has not completed the proper forms.

After Acceptance

Page proofs and corrections: Corresponding authors will receive electronic page proofs to check the copyedited and typeset article before publication. Portable document format (PDF) files of the typeset pages and support documents e.g., reprint order form) will be sent to the corresponding author by e-mail. Complete instructions will be provided with the e-mail for downloading and printing the files and for faxing the corrected pages to the publisher. Those authors without an e-mail address will receive traditional page proofs. It is the author's responsibility to ensure that there are no errors in the proofs. Changes that have been made to conform to journal style will stand if they do not alter the authors' meaning. Only the most critical changes to the accuracy of the content will be made. Changes that are stylistic or are a reworking of previously accepted material will be disallowed. The publisher reserves the right to deny any changes that do not affect the accuracy of the content. Authors may be charged for alterations to the proofs beyond those required to correct errors or to answer queries. Proofs must be checked carefully and corrections faxed within 24 to 48 hours of receipt, as requested in the cover letter accompanying the page proofs.

Electronic files should be submitted in a standard word processing format; Microsoft Word (or Corel WordPerfect) is preferred. Although conversions can be made from other word processing formats, the vagaries of the conversion process may introduce errors. Do not submit ASCII text files. Do not use automatic numbering or footnotes for references. The Journal does not assume responsibility for errors in the conversion of customized software, newly released software, and special characters. Authors preparing manuscripts on Macintosh computers should not use the Fast Save option.

Reprints: Authors will receive a reprint order form and a price list with the page proofs. Reprint requests should be faxed with the corrected proofs, if possible. Reprints are normally shipped 6 to 8 weeks after publication of the issue in which the item appears. Contact the Reprint Department, Lippincott Williams & Wilkins, 351 W. Camden Street, Baltimore, MD 21201; Fax: 410.528.4434; E-mail: reprints@lww.com with any questions.

Publisher's contact: Fax corrected page proofs, reprint order forms, and any other related material to Journal Production Editor, **The Clinical Journal of Pain**, 443-451-8199.

Manuscript Checklist (before submission)

- Cover letter with 3-4 suggested reviewers
- Title page
- Abstract
- References double-spaced in Journal style
- Corresponding author designated (in cover letter and on title page)
- E-mail address of corresponding author included in cover letter and on title page
- Permission to reproduce copyrighted materials or signed patient consent forms
- Acknowledgments listed for grants and technical support
- Authorship Responsibility, Financial Disclosure, and Copyright Transfer form signed by each author
- High-quality files of electronic art
- Appendix J- Permission for copyrighted questionnaires

Appendix I- Permission for Questionnaires

RE: Order Form for Department of Symptom Research Assessment Tools

symptomresearch [symptomresearch@mdanderson.org]

Sent: 29 August 2014 15:55

To: [Doughty Caitriona \(NHS FIFE\)](#)

Cc: [symptomresearch \[symptomresearch@mdanderson.org\]](#)

Attachments:

[BPI-SF English-24h Original.pdf \(61 KB\)](#)[\[Open as Web Page\]](#)

Hello

Caitriona,

I have attached the BPI as you requested. Please note that:

- Your use of the BPI is limited only to the study specified. To use the BPI in additional studies, you must reapply online at www.mdanderson.org/departments/prg > Symptom Assessment Tools > The Brief Pain Inventory (BPI).

- You are permitted to reproduce the copy of the BPI that is included with this e-mail. However, you must not remove the copyright notice.

- The BPI may not be modified in any way or translated into another language without the express written consent of the copyright holder; Charles S. Cleeland, PhD. Failure to comply may result in legal action. Permission to alter or translate the instrument may be obtained by contacting me at symptomresearch@mdanderson.org or by mail.

Please let me know if you have any questions. Thank you for your interest in the BPI.

Regards,

Mary Samad

Appendix J- IRAS protocol for Empirical Study

The relationship between self-compassion, self-esteem and pain in a chronic pain population



Caitríona Doughty (Principal Investigator)

Version 0.5 2015-05-24

Date: 24.05.15

Contents

Contacts	3
Introduction	4
Research Question	5
Study Sample	6
Inclusion and Exclusion Criteria	7
Consent process	6
Subject Withdrawal	6
Procedure	6
Data Collection	7
Analysis plan	10
Data Storage	10

Publication Policy	10
--------------------	----

References	17
------------	----

Contacts

Principal Investigator

*Caitríona Doughty (Protocol author)

Trainee Clinical Psychologist

Psychology Department

Lynebank Hospital

Halbeath Rd

DUNFERMLINE

KY11 4UW

Tel: 01383 565402

caitriona.doughty@nhs.net

Clinical Supervisor

Dr Laura Keith

Clinical Psychologist

Psychology Department

Lynebank Hospital

Halbeath Rd

DUNFERMLINE

KY11 4UW

Tel: 01383 565403

lkeith@nhs.net

Research Supervisor

Dr Paul G. Morris

Lecturer of Health Psychology

Clinical Psychology

Research Supervisor

Dr Nuno Ferreira

Lecturer

Clinical Psychology

School of Health in Social Science

Old Medical School,

Teviot Place

Edinburgh

EH8 9AG

School of Health in Social Science

Old Medical School

Teviot Place

Edinburgh

EH8 9AG

0131 651 3956

p.g.morris@ed.ac.uk

0131 650 3898

Nuno.ferreiro@ed.ac.uk

List of abbreviations

Scottish Intercollegiate Guidelines Network- SIGN; American College of Rheumatology- ACR; National Health Service- NHS; Self-Compassion Scale- SCS; Rosenberg Self-Esteem Scale- RSES; Brief Pain Inventory Short Form- BPISF; Quality of Life Scale-QOLS; Hospital Anxiety and Depression Scale- HADS.

Introduction

Chronic pain has been defined by Scottish Intercollegiate Guidelines Network (SIGN 136, 2013) as experiencing pain for at least 3 months that does not resolve with treatment. Approximately 20% of the adult European population have chronic pain (Breivik et al., 2006). Its management in the community remains generally unsatisfactory (van Hecke, Torrance & Smith, 2013). A specific chronic pain condition is fibromyalgia, which is diagnosed when an individual presents with widespread soft tissue pain (11 out of 18 paired tender points) lasting at least three months (Wolfe et al., 1990). Fibromyalgia is estimated to affect approximately 2-5% in Western Europe. Although fibromyalgia is the most common chronic widespread pain condition, it is often under diagnosed (Perrot, 2008). New criteria released by American College of Rheumatology (ACR, 2010) has proposed that focusing on the severity of characteristic fibromyalgia symptoms rather than tender point examinations may improve accuracy of diagnosis.

It is now widely accepted that psychological factors influence the relationship between chronic pain and outcomes such as disability and distress (Turk et al., 2008). Therefore, it may be beneficial in terms of managing fibromyalgia and other chronic pain conditions to explore possible psychosocial factors that contribute to the development of psychological distress in these conditions.

The association between pain and mental health symptoms in fibromyalgia were reported to be significantly higher than those in other chronic pain conditions (Gormsen et al., 2010; Taylor, Jason, & Jahn, 2003; Branco et al, 2010). It has been suggested that different mechanisms may be responsible for the development of mood disorders in fibromyalgia compared to other chronic pain conditions.

In particular, self-esteem has been linked to psychological adjustment in various chronic diseases (Chong et al, 2009), and how self-esteem is suspected to be a vulnerability factor in fibromyalgia (Johnson et al., 1997; Michielsen et al., 2006). Previous literature suggests that low self-esteem causes more negative affect for chronic disease patients than healthy populations (Bisschop et al., 2004). As self-esteem can be an important coping resource, one of the concerns raised by the combination of low self-esteem and chronic illness is an increased difficulty in preventing the negative affect from developing into a more severe case of depression. Research demonstrated that patients with chronic pain who had lower self-esteem reported more negative affect, less positive affect and greater symptom severity than individuals with higher self-esteem (Juth, Smyth, Santuzzi, 2008).

Despite the benefits of self-esteem, it has been suggested that high self-esteem is related to a number of negative outcomes including distortions in self-knowledge and increased aggression (Baumeister et al, 2003; Croker & Park, 2004). Reasons for this may be that self-esteem relies on meeting standards and favourable comparison with others (Neff, 2009). For this reason, research has started to focus on the construct of self-compassion which has been defined as the ability to be aware and open about one's flaws, while still being kind towards oneself.

Research on self-compassion is in its early stages, they have found that self-compassion is a significant predictor of increased psychological adjustment in people with chronic medical conditions (Wren, Somers, & Wright, 2011). Studies have also found that self-compassion correlates negatively with depression and anxiety (MacBeth & Gumley, 2012). It must be noted that much of the research on self-compassion has been conducted in non-clinical samples. Overall the literature appears to support the claim that self-compassion benefits people, especially when they experience failures or shortcomings. Research suggests that inclusion of positive and negative components of self-compassion would be beneficial in identifying

the 'active' components (Gilbert et al., 2011; van Dam et al., 2011). This study aims to investigate how self-compassion and self-esteem relate to pain outcomes in a chronic pain population, while also comparing their relative importance as mediators between pain intensity and outcomes of distress, disability and quality of life. It also aims to examine participants with fibromyalgia as a subgroup to investigate if there are differences within this population in relation to self-compassion and self-esteem as this group has been identified as having significantly high levels of distress and impaired functioning compared with other chronic pain patient groups (Gormsen, Rosenberg, Bach, & Jensen, 2010).

Research Question

Does self-compassion and self-esteem mediate between pain intensity and distress, functioning and quality of life?

How do the self-compassion subscales relate to pain outcomes?

Secondary Questions:

How does self-compassion and self-esteem relate to pain outcomes in a fibromyalgia subgroup compared to the overall chronic pain-group?

Study Sample

Individuals will be recruited through NHS Fife Pain Management Service, NHS Fife Rheumatic Diseases Unit and chronic pain support groups. All documents will each be marked with a unique participant number which will not identify the participant personally, ensuring their anonymity.

Patient Inclusion and Exclusion criteria:

Inclusion

- * Must be at least 18 years of age
- * Able to speak, read and write fluently in English
- * Experienced chronic pain in any location of the body for a minimum of 3 months in line with the SIGN Guidelines (2013).

Exclusion

*Individual is suffering from malignant or terminal condition such as cancer.

*Individual has any of the below

-Intellectual impairment (e.g. Learning Disability, Alzheimers' Disease, Brain Injury)

-Severe mental illness

-Primary alcohol misuse

-Substance misuse

Number of participants

Given that no previous study has investigated the relationship between these variables in a chronic pain sample, the projected effect size for the current study has been conservatively estimated at a moderate level of $r^2 = 0.15$. Assuming a medium-size relationship between the predictors and the outcomes, $\alpha = .05$ and $\beta = .20$, the formula $N \geq 50 + 8m$ (where m is the number of predictors) can be used for testing the regression (Tabachnick & Fidell, 2007).

After calculating a G*power analysis, it was indicated that the minimum number of participants required for this study is 80. In order to carry out the appropriate statistics for this study with sufficient power, it is estimated that the required sample number is at least 120 participants.

Previous research has indicated that the response rate of postal survey studies among people with chronic pain range between 43-50% (Gunn et al., 2012). Given that the principal researcher aims to access up to 3 potential sources of recruitment, it is possible that at least 480 questionnaires could be circulated, which would guarantee the required sample size.

Consent Process

Participants will be recruited from NHS Fife Pain Management Service and the NHS Fife Rheumatoid Disease Unit. This will be a single site study in that participants will only be recruited the NHS Fife area. It is anticipated that the majority of participants will be identified by NHS Fife clinical staff (Clinical Psychology, Nurse Specialists) through their clinical role with participants. Clinicians will be asked to approach any patients they believe fulfil the inclusion/exclusion criteria and to provide them with the participant information and consent sheet. Potential participants will not be identified by any personal or clinical information.

Consent will be implied if the individual completes the questionnaires and posts them back to the address on the pre-paid envelope

Participant Withdrawal

The information provided by the participant for the research study will not be identifiable. Therefore, it will not be possible to remove their responses from the study after they have posted them back to the researcher. This will be made explicitly clear to the potential participants.

Procedure

Taking part in the study involves completing a number of questionnaires, which should take approximately thirty minutes to complete. The participant will also be asked to provide some very basic details e.g. gender, age, type of diagnosis which will not identify them personally. If they agree to take part in the study, they will be asked to post them back using the pre-paid envelope included.

Data collection

The study is a cross-sectional design. Attendees at the Fife Pain Management Service (NHS), the Rheumatic Disease Unit (NHS) and a voluntary sector chronic pain support groups (not affiliated with the NHS) will be recruited. They will be given a questionnaire pack that contains information sheets, a pre-paid stamped-addressed envelope and six validated questionnaires of pain, self-compassion, self-esteem, distress, disability and quality of life.

Permission has been granted by the primary authors for use of all of the questionnaires.

Self-Compassion Scale (SCS; Neff, 2003) is a 26 item scale that assesses levels of self-compassion. There are three factors of positive self-compassion: self-kindness, common humanity and mindfulness and three factors that focus on a lack of self-compassion and negative self-evaluation: self-judgement, isolation and over-identification. Participants indicate how often they engage in these ways of self-relating on a Likert scale 1-5. The scale has good reliability with Cronbach's alpha ranging from .72 to .83 (van Dam, Sheppard, Forsyth, & Earleywine, 2011).

Rosenberg Self Esteem Scale (RSES; Rosenberg, 1965) measures a global feeling of self-worth or self-acceptance. It is rated on a 4 point scale from 1, strongly agree to 4, strongly disagree. The ten statements are designed equally to be positive and negative. The total score ranges from 0 to 30. The scale has good reliability with a Cronbach's alpha of 0.89 (Michielsen, van Houdenhove, Leirs, Vandenbrock, & Onghena, 2006).

Brief Pain Inventory Short Form (BPI-SF; Cleeland, 1991) contains a total of 15 items and assesses for the presence of pain, pain intensity and functional interference from pain. It also catalogues the types of pain medications being used; the percentage of pain relief obtained from medications and assesses the distribution of pain via a body map. The BPI-SF is widely used in clinical trials for pain and is one of the instruments recommended by the initiative on methods measurement and pain assessment in clinical trials group (Dwarkin et al., 2005). Internal consistency for the Pain Severity Score and for the Interference scale has been reported as ranging from 0.85 and 0.88 respectively (jensen et al., 2004). Construct validity of the BPI has been supported for the generic assessment of pain as well as specifically for low back pain and rheumatoid arthritis (Keller et al., 2004).

Pain Disability Questionnaire (PDI; Anagnostis, Gatchel, & Mayer, 2004) incorporates a disability-related psychosocial component in addition to a physical functioning component related to pain. It yields a total functioning disability score ranging from 0 (optimal function) to 150 (total disability), using 11 point Likert scale (Gatchel et al., 2006). The PDQ has good reliability with a Cronbach's alpha of .96. The construct related validity of the PDQ was found to be excellent as it correlated well to both the MVAS (0.65-0.81) and Owesstry (0.55-0.80) (Anagnostis et al, 2004).

Quality of Life Scale (QOLS; Burckhardt, Woods, Schultz, & Ziebarth, 2003) is a 16 item questionnaire that includes various aspects of life such as physical and material well-being, relationships with other people, social, community and civic activities, personal development and fulfilment, recreation and independence. Each item was scored 0-7 and the total score ranges 16-112. A higher total score shows a higher QOL. This scale has shown good internal consistency ($\alpha = 0.88$), validity and test-retest reliability (Pedro et al, 2014).

The Hospital Anxiety and Depression Scale (HADS; Zigmond and Snaith, 1983) is widely used for measuring anxiety and depression in clinical populations with physical illness. It contains 14 items and gives an overall measure of emotional distress as well as depression and anxiety subscale scores. It has shown good internal consistency ($\alpha = 0.90-0.93$), validity and test-retest reliability (Moorey et al., 1991; Snaith and Zigmond, 1994; Bjelland et al., 2002). In this study, a total score was used to represent emotional distress.

Table 1: Research Timescale

Feb	Mar	A pr	M ay	J u n	J u l	A u g	S e p	O ct	N ov	D ec	J a n	Feb	Ma r	Apr	May
Ethi cs															
	Data Collecti on	*	*	*	*	*	*	*	*	*	*	End of Data Collecti on			
													Data Analysi s		
													Wri te Up	Wri te Up	
															Submissi on

Analysis Plan

Preliminary analyses will be performed to check for missing data, normality and adequacy for mediation analysis. A set of bivariate Pearson's correlation analyses will explore the relations between the proposed mediators of pain and pain outcomes (distress, function and quality of life). Of additional interest will be to use multiple regression to investigate the extent that the subscales of the SCS make individual contributions to predictive validity between pain and pain outcomes.

Multiple mediational analysis using the bootstrapping approach as described by Preacher and Hayes (2008) and Hayes (2009), will be used to investigate the relative importance that self-compassion and self-esteem have in the mediation of the relation between pain experiences (intensity) and pain outcomes (physical and emotional). To evaluate the magnitude of indirect effects, the product of co-efficients approach (Bishop, Fienberg, & Holland, 2008) will be used to calculate standard errors of the indirect effects.

To answer the second question of the study, a series of t-tests will be used to detect differences between the Chronic Pain and the Fibromyalgia.

Data Storage

All data will be stored in a manner consistent with NHS Fife guidance and policy on confidentiality and IT security, and in keeping with data protection legislation. Any of the data removed from NHS systems (e.g. for academic project work) will already be fully anonymised therefore, patient's anonymity will be preserved.

Publishing Outcomes

A report of the findings will be provided to Edinburgh University, the NHS Fife Psychology Department and submitted for publication in a peer reviewed journal.

References

- Anagnostis, C. Gatchel, R., & Mayer, T. (2004). The pain disability questionnaire: A new psychometrically sound measure for chronic musculoskeletal disorders. *Health Services Research, 29*, 2290-2302.
- Baumeister, R.F. (2003). Does high self-esteem cause better performance, interpersonal success, happiness, or healthier lifestyles? *Psychological Science in the Public Interest, 4*, 1-45.
- Bishop, Yvonne M., Stephen E. Fienberg, and Paul W. Holland. *Discrete multivariate analysis: theory and practice*. Springer, 2007.
- Bisschop, M.I. (2004). Chronic diseases and depression: the modifying role of psychosocial resources. *Social Science & Medicine, 59*, 721-733.
- Branco, J.C. (2010). Prevalence of fibromyalgia: A survey in five European countries. *Seminars in Arthritis & Rheumatism, 39*, 448-453.
- Breivik, H., Collett, B., Ventafridda, V., Cohen, R. & Gallacher, D. (2006). Survey of chronic pain in Europe: Prevalence, impact on daily life and treatment. *European Journal of Pain, 10*, 287-293.
- Burckhardt, C.S. & Anderson, K.L. (2003). The quality of life scale (qols): Reliability, validity and utilisation. *Health and Quality of Life Outcomes, 60*, 1-7.
- Cleeland CS. (1991). The Brief Pain Inventory-Short Form.
- Crocker, J. & Park, L. (2004). The costly pursuit of self-esteem. *Psychological Bulletin, 130*, 392-414.
- Gormsen, L., Rosenberg, R., Bach, F.W. & Jensen, T.S. (2010). Depression, anxiety, health-related quality of life and pain in patients with chronic fibromyalgia and neuropathic pain. *European Journal of Pain, 14*, 127-133.
- Johnson, M., Paananen, M.L., Rahainantti, P. & Hannonen, P. (1997). Depressed fibromyalgia patients are equipped with an emphatic competence dependent self-esteem. *Clinical rheumatology, 16*, 578-584.

- Juth, V, Smyth, J.M., & Santuzzi, A.M. (2010). How do you feel? Self-esteem predicts affect, stress, social interaction, and symptom severity during daily life in patients with chronic illness. *Journal of Health Psychology, 13*, 884-894.
- MacBeth, A. & Gumley, A. (2012). Exploring compassion: A meta-analysis of the association between self-compassion and psychopathology. *Clinical Psychology Review, 32*, 545-552.
- Michielsen, H.J., van Houdenhove, B., Leirs, I., Vandenbroeck, A., & Onghena, P. (2006). Depression, attribution style and self-esteem in chronic fatigue syndrome and fibromyalgia patients: Is there a link? *Clinical Rheumatology, 25*, 183-188.
- Neff, K. (2009). The role of self-compassion in development: A healthier way to relate to oneself. *Human Development, 52*, 211-214.
- Neff, K. (2003). Development and validation of a scale to measure self-compassion. *Self and Identity, 2*, 223-250.
- Perrot, S. (2008). Fibromyalgia: Harmonising science with clinical practice considerations. *Pain Practice, 8*, 177-189.
- Preacher, K.J. & Hayes, A. (2008). Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behaviour Research Methods, 40*, 879-891.
- Rosenberg, Morris. (1965). "Rosenberg Self-Esteem Scale (RSE)." *Acceptance and Commitment Therapy. Measures Package*: 61.
- Taylor, R.R. Jason, L., & Jahn, S. (2003). Chronic fatigue and sociodemographic characteristics as predictors of psychiatric disorders in a community-based sample. *Psychosomatic Medicine, 65*, 896-901.
- Turk, D.C., Swanson, K.S., Tunks, E.R. (2008). Psychological approaches in the treatment of chronic pain patients- when pill, scalpels and needles are not enough. *Canadian Journal of Psychiatry, 53*, 213-223.

Van Dam, N., Sheppard, S., Forsyth, J.P. (2011). Self-compassion is a better predictor than mindfulness of symptom severity and quality of life in mixed anxiety and depression. *Journal of Anxiety Disorders*, 25, 123-130.

Van Hecke, O., Torrance, N., Smith, B.H. (2013). Chronic pain epidemiology and its clinical relevance. *British Journal of Anaesthesia*, 111, 13-18.

Wolfe, F. (2010). The American college of rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care and Research*, 62, 600-610.

Wolfe, F. (1990). The American college of rheumatology 1990 criteria for the classification of fibromyalgia. *Arthritis and Rheumatism*, 33, 160-172.

Wren, A., Somer, T. Wright, M. (2012). Self-compassion in patients with persistent musculoskeletal pain: Relationship of self-compassion to adjustment to persistent pain. *Journal of Pain and Symptom Management*, 43, 759-770.

Zigmond, A.S., & Snaith, R.P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, 67, 361-370.

Appendix K-Ethical Approval Letters

South East Scotland REC 01

Waverley Gate
2 - 4 Waterloo Place
Edinburgh
EH1 3EJ

Telephone: 0131 465 547

26 March 2015

Ms Caitriona Doughty
Trainee Clinical Psychologist
NHS Fife
Lynebank Hospital
Halbeath Road
Dunfermline
KY11 4UW

Dear Ms Doughty

Study title: The Role of Self Compassion and Self-Esteem in the Relationship between Chronic Pain and outcomes of Psychological Distress, Functioning, and Quality of Life
REC reference: 15/SS/0035
IRAS project ID: 147820

Thank you for your letter of 24 March 2015. I can confirm the REC has received the documents listed below and that these comply with the approval conditions detailed in our letter dated 17 March 2015

Documents received

The documents received were as follows:

<i>Document</i>	<i>Version</i>	<i>Date</i>
IRAS Checklist XML [Checklist_24032015]		24 March 2015
Participant information sheet (PIS)	6	23 March 2015
Validated questionnaire [Brief Pain Inventory SF]	V2	23 March 2015
Validated questionnaire [Self Compassion Scale]	2	23 March 2015
Validated questionnaire [Pain Disability Questionnaire]	V3	26 March 2015
Validated questionnaire [Quality of Life]	V2	23 March 2015

Approved documents

The final list of approved documentation for the study is therefore as follows:

<i>Document</i>	<i>Version</i>	<i>Date</i>
Non-validated questionnaire [caitrona doughty demographic questionnaire]		
Participant information sheet (PIS)	6	23 March 2015
REC Application Form [REC_Form_16022015]		16 February 2015
Research protocol or project proposal	Version 0.5	13 February 2015
Summary CV for Chief Investigator (CI)		
Summary CV for supervisor (student research)		
Validated questionnaire [Rosenberg Self Esteem Scale]		
Validated questionnaire [Hospital Anxiety and Depression Scale]		
Validated questionnaire [Brief Pain Inventory SF]	V2	23 March 2015
Validated questionnaire [Self Compassion Scale]	2	23 March 2015
Validated questionnaire [Pain Disability Questionnaire]	V3	26 March 2015
Validated questionnaire [Quality of Life]	V2	23 March 2015

You should ensure that the sponsor has a copy of the final documentation for the study. It is the sponsor's responsibility to ensure that the documentation is made available to R&D offices at all participating sites.

15/SS/0035	Please quote this number on all correspondence
-------------------	---

Yours sincerely



Sandra Wyllie
REC Manager

E-mail: sandra.wyllie@nhslothian.scot.nhs.uk

Copy to: Ms Caitriona Doughty, NHS Fife
Mrs Jo-Anne Robertson
Ms Aileen Yell, NHS Fife

Medical Director

Hayfield House
Hayfield Road
KIRKCALDY
KY2 5AH



Ms Caitriona Doughty
Trainee Clinical Psychologist
Psychology Dept
Lynebank Hospital
DUNFERMLINE

Our Ref 31 March 2015
15-001 15/SS/0035

Enquiries to Aileen Yell
E-mail aileenyell@nhs.net
Telephone 01383 623623 Ext 20940
Website www.nhsfife.org

Dear Ms Doughty

Project Title: The role of self compassion and self-esteem in the relationship between chronic pain and outcomes of psychological distress, functioning and quality of life

Thank you for your application to carry out the above project. Your project documentation (detailed below) has been reviewed for resource and financial implications for NHS Fife and I am happy to inform you that NHS permission for the above research has been granted on the basis described in the application form, protocol and supporting documentation. The documents reviewed were:

Document	Version	Date
Protocol	0.5	13 February 2015
REC provisional favourable opinion letter		17 March 2015
Participant Information Sheet	6	23 March 2015
REC final favourable opinion letter		26 March 2015
IRAS R&D Form	3.5	27 March 2015
IRAS SSI Form	3.5	27 March 2015

The terms of the approval state that you are the Principal Investigator authorised to undertake this study within NHS Fife, with assistance from Jane Timperley in the Chronic Pain Management Team.


I note that the favourable ethical opinion applies to all NHS sites taking part in the study therefore no separate Site Specific Review is required in this case. The sponsors for this study are University of Edinburgh.

Details of our participation in studies will be included in annual returns we are expected to complete as part of our agreement with the Chief Scientist Office. Regular reports of the study require to be submitted. Your first report should be submitted to Dr A Wood, R&D Manager, R&D Department, Queen Margaret Hospital, Whitefield Rd, Dunfermline, KY12 0SU (Amanda.wood3@nhs.net) in 12 months time and subsequently at yearly intervals until the work is completed. A Lay Summary will also be required upon completion of the project.

In addition, approval is granted subject to the following conditions:-

All research activity must comply with the standards detailed in the Research Governance Framework for Health & Community Care (<http://www.cso.scot.nhs.uk/publications/resgov/resgov.htm>), health & safety regulations, data

⁶ NHS Fife was awarded the Carbon Trust Standard in February 2010



protection principles, other appropriate statutory legislation and in accordance with Good Clinical Practice (GCP).

Any amendments which may subsequently be made to the study should also be notified to Aileen Yell, Research Governance Officer (aileenyell@nhs.net), as well as the appropriate regulatory authorities. Notification should also be given of any new research team members post approval and/or any changes to the status of the project.

This organisation is required to monitor research to ensure compliance with the Research Governance Framework and other legal and regulatory requirements. This is achieved by random audit of research. You will be required to assist with and provide information in regard to monitoring and study outcomes (including providing recruitment figures to the R&D office as and when required). Please send a copy of all Monitoring Reports submitted to the REC to Allyson.bailey@nhs.net


As custodian of the information collated during this research project you are responsible for ensuring the security of all personal information collected in line with NHS Scotland IT Security Policies, until the destruction of this data.

Permission is only granted for the activities for which a favourable opinion has been given by the REC (and which have been authorised by the MHRA where appropriate).

The research sponsor or the Chief Investigator or local Principal Investigator at a research site may take appropriate urgent safety measures in order to protect research participants against any immediate hazard to their health or safety. The R&D office (aileenyell@nhs.net) should be notified that such measures have been taken. The notification should also include the reasons why the measures were taken and the plan for further action. The R&D office should be notified within the same time frame of notifying the REC and any other regulatory bodies.

I would like to wish you every success with your study and look forward to receiving a summary of the findings for dissemination once the project is complete.

Yours sincerely



DR BRIAN MONTGOMERY
Medical Director
NHS Fife

Cc : Aileen Yell, Research Governance Officer, NHS Fife, Queen Margaret Hospital, Dunfermline

University Hospitals Division

Queen's Medical Research Institute
47 Little France Crescent, Edinburgh, EH16 4TJ

FM/CK/approval

30 July 2015

Dr Evelyn Janetta
NHS Lothian
Pain Management Programme
Astley Ainslie Hospital
133 Grange Loan
Edinburgh
EH10 2HL



Research & Development
Room E1.12
Tel: 0131 242 3330

Email:
R&DOffice@nhslothian.scot.nhs.uk

Director: Professor David E Newby

Dear Dr Janetta

Lothian R&D Project No: 2015/0279

Title of Research: The role of self compassion and self-esteem in the relationship between chronic pain and outcomes of psychological distress, functioning and quality of life

REC No: 15/SS/0035

Participant Information Sheet:

Version 6.0, 20 March 2015

Consent Form:

N/A

Protocol: Version 5.0, 13 February 2015

I am pleased to inform you that this study has been approved for NHS Lothian and you may proceed with your research, subject to the conditions below. This letter provides Site Specific approval for NHS Lothian.

Please note that the NHS Lothian R&D Office must be informed if there are any changes to the study such as amendments to the protocol, recruitment, funding, personnel or resource input required of NHS Lothian.

Substantial amendments to the protocol will require approval from the ethics committee which approved your study and the MHRA where applicable.

Please inform this office when recruitment has closed and when the study has been completed.

I wish you every success with your study.

Yours sincerely

Fiona McArdle

Ms Fiona McArdle
Deputy R&D Director

Cc: Dr Caroline Cochrane, Consultant Clinical Psychologist, Lead, Astley Ainslie Hospital
Ms Caitriona Doughty, Trainee Clinical Psychologist, Lynebank Hospital
Ms Jenny Fleming, General Manager, Western General Hospital



SCHOOL of HEALTH IN SOCIAL SCIENCE
CLINICAL AND HEALTH PSYCHOLOGY

The University of Edinburgh
Medical School
Doorway 6, Teviot Place
Edinburgh EH8 9AG

Telephone 0131 651 3969
Fax 0131 650 3891
Email submitting.ethics@ed.ac.uk

Caitriona Doughty
Trainee Clinical Psychologist
Dept of Psychology
Lynebank Hospital
Halbeath Road
Dunfermline
KY11 4UW

13 April 2016

Dear Caitriona,

Application for Level 1 Ethical Approval

Reference: CLIN266

Project Title: Self Compassion, Self Esteem and Chronic Pain

Academic Supervisors: Paul Morris / Nuno Ferreira

Thank you for submitting the above research project for review by the Department of Clinical and Health Psychology Ethics Research Panel. I can confirm that the submission has been independently reviewed and was approved on the 30th March 2016.

Should there be any change to the research protocol it is important that you alert us to this as this may necessitate further review.

Yours sincerely,

Kirsty Gardner
Administrator
Clinical Psychology



The relationship between self-compassion, self-esteem and pain in a chronic pain population

We would like to invite you to take part in our research study. Before you decide, we would like you to understand why the research is being done and what it would involve for you. If you have any questions about this study not answered in this information sheet, you can contact the researcher, Caitríona Doughty.

What is the purpose of the study?

The purpose of the study is to improve understanding of the psychological factors (e.g. self-compassion and self-esteem) that contribute to the development of psychological distress and disability in chronic pain conditions. Clinical psychologists conduct research to contribute to the evidence about factors affecting mental health, in order to improve therapeutic approaches such as talking therapies. This study is being conducted as part fulfilment of the researcher's academic requirements of the Clinical Psychology Doctorate at University of Edinburgh. No one is being paid for your participation in this study.

Why have I been invited?

You have been contacted because you are an individual experiencing chronic pain for over 3 months, who has been attending any of the following service: NHS Fife Pain Management Service, NHS Fife Rheumatic Diseases Unit, Chronic Pain Self Management Group in Fife, NHS Lothian Pain Management Programme or NHS Lothian Rheumatology.

Do I have to take part?

No, you do not have to take part and you do not have to provide a reason. Your decision will not affect the standard of care or support you receive. You will be asked by a health care provider or group facilitator if you would like a questionnaire pack for this study. If you agree to this, you can then take this home with you and decide whether you would like to participate. This study will be accepting response questionnaires from February 2015 to February 2016. **If you complete and return the attached questionnaires, this will be taken as your consent to participate.** Written consent is not being gathered in this study in order to maintain anonymity.

What am I being asked to do?

Taking part in this research involves completing a number of questionnaires, which should take approximately 35 minutes to complete. You will also be asked to provide some very basic details, which will not identify you personally in any way e.g. gender. If you agree to take part in this study you are asked **to post them back using the pre-paid envelope included**. This will be the only time information about you, or from you, will be required for this study and it will be the end of your participation in the study.

What are the possible disadvantages and risks of taking part?

The risks or disadvantages identified are minimal when taking part in this study. Completion of the questionnaires may be inconvenient and you may find some of the questions upsetting. You can stop completing the questionnaires at any time. You should speak to your GP if your distress continues. Listed below are a number of resources that you can access if you feel distressed. Included in this pack is a leaflet providing contact details for a confidential helpline service known as Breathing Space.

Moodcafe- The Well-being Website

Website Address: www.moodcafe.co.uk

Pain Association Scotland- Self Management for Chronic Pain

Contact telephone number: 0800 783 6059

What are the possible benefits of taking part?

While there are no direct benefits to taking part in this study, you may enjoy contributing to research that is intended to improve future patient care by providing a broader understanding of the factors that influence pain outcomes in a chronic pain population.

What if there is a problem?

If you have a concern or complaint about any aspect of this study, you should contact the researcher who will do her best to answer your questions (Caitríona Doughty, 01383 565402). If you would like to speak to an independent adviser who can answer questions about the study but is not directly involved in running the research you can contact - Tara Graham (Research & Service Development Psychologist), Department of Clinical Psychology Stratheden Hospital, Cupar, KY15 5RR. (01334 696336).

If you remain unhappy and wish to complain formally, you can contact Dr Katherine Cheshire, Head of Clinical Psychology Department, NHS Fife (01383 565403), or Paul Morris, Academic Tutor for the study (0131 6513956). You may also follow the NHS complaints route by contacting the Patient Relations Department office, Room 104, Hayfield House, Hayfield Road, Kirkcaldy, KY2 5AH, (01592 648153) or email patientrelations.fife@nhs.net.

What will happen if I don't want to carry on with the study?

The information you provide for the research study will not identify you. Therefore, once you have returned the questionnaires it will not be possible to remove your responses from the study should you change your mind.

Will my taking part in this study be kept confidential?

Yes. We will follow ethical and legal practice and all information about you will be handled in confidence. Information you provide for this study will not contain any personally identifiable information and therefore your participation will be confidential.

What will happen to the results of the research study?

A report of the findings will be provided to Edinburgh University, the NHS Fife Psychology Department and submitted for publication in a peer reviewed journal. If you require a copy of the report, please contact the researcher. Participants will not be identified in any way.

Who has reviewed the study?

The South East Scotland Research Ethics Committee 01, which has responsibility for scrutinising all proposals for medical research on humans in Tayside, has examined the proposal and has raised no objections from the point of view of medical ethics.

Thank you

Appendix M- Demographic Information Sheet



The Relationship Between Self-Compassion, Self-Esteem and Pain in a Chronic Pain Population

Demographic Information

Please answer the following questions

Recruitment site (researcher to complete prior to sending out packs)	
Gender	
Age	
How long have you experienced chronic pain	
Please list any diagnosis you have been given in relation to your chronic pain	
Was this diagnosis provided by your GP	YES <input type="checkbox"/> NO <input type="checkbox"/>
Other, please specify	

Thank you.